



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04090187.8

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04090187.8
Demande no:

Anmeldetag:
Date of filing: 10.05.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Epigenomics AG
Kastanienallee 24
10435 Berlin
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Methods and nucleic acids for the analysis of CpG dinucleotide methylation status
associated with the development of prostate cancer

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

EP/10.02.04/EP 04090040
EP/01.12.03/EP 03090414

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12Q1/68

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PL PT RO SE SI SK TR LI

METHODS AND NUCLEIC ACIDS FOR THE ANALYSIS OF CpG DINUCLEOTIDE
METHYLATION STATUS ASSOCIATED WITH THE DEVELOPMENT OF PROSTATE
CANCER. EPO-BERLIN

10-05-2004

FIELD OF THE INVENTION

The present invention relates to human DNA sequences that exhibit altered methylation patterns (hypermethylation or hypomethylation) in cancer patients. Particular embodiments of the invention provide highly accurate methods for detection and differentiation of prostate carcinomas.

BACKGROUND

Correlation of aberrant DNA methylation with cancer. Aberrant DNA methylation within CpG 'islands' is characterized by hyper- or hypomethylation of CpG dinucleotide sequences leading to abrogation or overexpression of a broad spectrum of genes, and is among the earliest and most common alterations found in, and correlated with human malignancies. Additionally, abnormal methylation has been shown to occur in CpG-rich regulatory elements in intronic and coding parts of genes for certain tumors. In colon cancer, aberrant DNA methylation constitutes one of the most prominent alterations and inactivates many tumor suppressor genes including, *inter alia*, p14ARF, p16INK4a, THBS1, MINT2, and MINT31 and DNA mismatch repair genes such as hMLH1.

Aside from the specific hypermethylation of tumor suppressor genes, an overall hypomethylation of DNA can be observed in tumor cells. This decrease in global methylation can be detected early, far before the development of frank tumor formation. A correlation between hypomethylation and increased gene expression has been determined for many oncogenes.

Prostate cancer. The prostate is a male sex accessory gland, comprising about 30 to 50 branched glands. It is surrounded by a fibroelastic capsule that separates the gland into discrete lobes. The central zone of the organ is composed of pseudo stratified epithelium, the peripheral zone comprises the bulk of the organ and the two tissue types are separated by a transitional zone.

Benign prostate hypertrophy is present in about 50% of men aged 50 or above, and in 95% of men aged 75 or above. Prostate

cancer is a significant health care problem in Western countries with an incidence of 180 per 100,000 in the United States in 1999 (*Cancer J. Clin.*, 49:8, 1999).

Diagnosis and prognosis of prostate cancer; deficiencies of prior art approaches. Different screening strategies have been employed with at least some degree of success to improve early detection of prostate cancer, including determination of levels of prostate specific antigen ("PSA") and digital rectal examination. If a prostate carcinoma is suspected in a patient, diagnosis of cancer is confirmed or excluded by the histological and cytological analysis of biopsy samples for features associated with malignant transformation. The zone of origin of a prostatic cell proliferative disorder is currently determined by the 'PSA density.' PSA density is determined by dividing the weight of the prostate (as estimated by transrectal ultrasound) by the prostate specific antigen levels of the patient. Levels of over 15% percent are considered as indicative of prostate cancer and grounds for a biopsy. The biopsy, in turn, is used for histological and cytological analysis to determine the zone of origin.

However, using routine histological examination, it is often difficult to distinguish benign hyperplasia of the prostate from early stages of prostate carcinoma, even if an adequate biopsy is obtained (McNeal J. E. et al., *Hum. Pathol.* 2001, 32:441-6). Furthermore, small or otherwise insufficient biopsy samples often impede the analysis.

Molecular markers would offer the advantage that they could be used to efficiently analyze even very small tissue samples, and samples whose tissue architecture has not been maintained. Within the last decade, numerous genes have been studied with respect to differential expression among benign hyperplasia of the prostate and different grades of prostate cancer.

However, no single marker has as yet been shown to be sufficient for the diagnosis of prostate tumors in a clinical setting.

Alternatively, high-dimensional mRNA based approaches may, in particular instances, provide a means to distinguish between different tumor types and benign and malignant lesions. However, application of such approaches as a routine diagnostic

tool in a clinical environment is impeded and substantially limited by the extreme instability of mRNA, the rapidly occurring expression changes following certain triggers (e.g., sample collection), and, most importantly, by the large amount of mRNA needed for analysis which often cannot be obtained from a routine biopsy (see, e.g., Lipshutz, R. J. et al., *Nature Genetics* 21:20-24, 1999; Bowtell, D. D. L. *Nature Genetics Suppl.* 21:25-32, 1999).

The GSTP1 gene. The core promoter region of the Glutathione S-Transferase P gene (GSTP1; accession no. NM_000852) has been shown to be hypermethylated in prostate tumor tissue. The glutathione S-transferase pi enzyme is involved in the detoxification of electrophilic carcinogens, and impaired or decreased levels of enzymatic activity (GSTPi impairment) have been associated with the development of neoplasms, particularly in the prostate. Mechanisms of GSTPi impairment include mutation (the GSTP*B allele has been associated with a higher risk of cancer) and methylation.

Prior art GSTP1 studies. Lee et al., in United States Patent No 5,552,277, disclosed that the expression of the glutathione-S-transferase (GST) Pi gene was downregulated in a significant proportion of prostate carcinomas. Moreover, by means of restriction enzyme analysis they were able to show that the promoter region of the of the GSTPi gene was upmethyalted (hypermethylated) in prostate carcinomas as opposed to normal prostate and leukocyte tissue. However, due to the limited and imprecise nature of the analysis technique used (HpaIII digestion, followed by Southern blotting) the exact number and position of the methylated CG dinucleotides were not characterized.

Douglas et al. (WO9955905) used a method comprising bisulfite treatment, followed by methylation specific PCR to show that prostate carcinoma-specific GSTPi hypermethylation was localized to the core promoter regions, and localized a number of CpG positions that had not been characterised by Lee et al.

Herman and Baylin (United States Patent No. 6,017,704) describe the use of methylation specific primers for methylation analysis, and describe a particular primer pair

suitable for the analysis of the corresponding methylated GSTPi promoter sequence.

However, with respect to the use of GSTPi markers, the prior art is limited with respect to the number of GSTPi promoter CpG sequences that have been characterized for differential methylation status. Moreover, there are no disclosures, suggestions or teachings in the prior art of how such markers could be used to distinguish among benign hyperplasia of the prostate and different grades of prostate cancer.

Aberrant genetic methylation has also been observed in several other genes including AR, p16 (CDKN2a/INK4a), CD44, CDH1. Genome wide hypomethylation for example of the LINE-1 repetitive element has also been associated with tumor progression (Santourlidis S ,Florl A ,Ackermann R ,Wirtz HC ,Schulz WA 'High frequency of alterations in DNA methylation in adenocarcinoma of the prostate.' Prostate 1999 May 15;39(3):166-74).

However, use of these genes as alternative or supplemental diagnostic, or otherwise clinically useful markers in a commercial setting has not been enabled. The application of differentially methylated genes to clinically utilizable platforms requires much further investigation into the sensitivity and specificity of the genes. For example, in the case of the gene CD44, a known metastasis suppressor, downregulation was associated with hypermethylation. However the use of this gene as a commercially available marker was not enabled as it was also methylated in normal tissues. See Vis AN Oomen M Schroder FH van der Kwast TH 'Feasibility of assessment of promoter methylation of the CD44 gene in serum of prostate cancer patients.' Mol Urol. 2001 Winter;5(4):199-203.

Development of medical tests. Two key evaluative measures of any medical screening or diagnostic test are its sensitivity and specificity, which measure how well the test performs to accurately detect all affected individuals without exception, and without falsely including individuals who do not have the target disease (predictive value). Historically, many diagnostic tests have been criticized due to poor sensitivity and specificity.

A true positive (TP) result is where the test is positive and the condition is present. A false positive (FP) result is where the test is positive but the condition is not present. A true negative (TN) result is where the test is negative and the condition is not present. A false negative (FN) result is where the test is negative but the condition is not present.

Sensitivity = $TP / (TP + FN)$

Specificity = $TN / (FP + TN)$

Predictive value = $TP / (TP + FP)$

Sensitivity is a measure of a test's ability to correctly detect the target disease in an individual being tested. A test having poor sensitivity produces a high rate of false negatives, i.e., individuals who have the disease but are falsely identified as being free of that particular disease. The potential danger of a false negative is that the diseased individual will remain undiagnosed and untreated for some period of time, during which the disease may progress to a later stage wherein treatments, if any, may be less effective. An example of a test that has low sensitivity is a protein-based blood test for HIV. This type of test exhibits poor sensitivity because it fails to detect the presence of the virus until the disease is well established and the virus has invaded the bloodstream in substantial numbers. In contrast, an example of a test that has high sensitivity is viral-load detection using the polymerase chain reaction (PCR). High sensitivity is achieved because this type of test can detect very small quantities of the virus. High sensitivity is particularly important when the consequences of missing a diagnosis are high.

Specificity, on the other hand, is a measure of a test's ability to identify accurately patients who are free of the disease state. A test having poor specificity produces a high rate of false positives, i.e., individuals who are falsely identified as having the disease. A drawback of false positives is that they force patients to undergo unnecessary medical procedures treatments with their attendant risks, emotional and

financial stresses, and which could have adverse effects on the patient's health. A feature of diseases which makes it difficult to develop diagnostic tests with high specificity is that disease mechanisms, particularly in cancer, often involve a plurality of genes and proteins. Additionally, certain proteins may be elevated for reasons unrelated to a disease state. An example of a test that has high specificity is a gene-based test that can detect a p53 mutation. Specificity is important when the cost or risk associated with further diagnostic procedures or further medical intervention are very high.

The PSA blood test has a sensitivity of 73%, specificity of 60% and predictive value of 31.5%. PSA sensitivity and specificity can be improved but involve tradeoffs. PSA sensitivity can be improved by adjusting the "normal" PSA level to a lower value for younger men or by following serum PSA values in an individual patient over time (PSA velocity). Both methods will increase the number of cancers detected, but they also increase the number of men undergoing biopsy. Conversely, specificity can be improved by using higher "normal" PSA levels for older men, by using the free-to-total PSA ratio, or by adjusting the normal value according to the size of the prostate. These three methods decrease the number of unnecessary biopsies, but they increase the risk that some cancers will be missed.

It can therefore be seen that there exists a need for a means of prostate cancer diagnosis with improved sensitivity, specificity and/or predictive value.

Sensitivity and specificity of quantitative methylation-specific polymerase chain reaction (QMSP) assay alone (without histological analysis) in prostate cancer analysis of needle biopsies has ranged from 30% sensitivity and 100% specificity to 89% sensitivity and 64% specificity (Harden et. al. J Natl Cancer Inst 2003; 95: 1634-1637). However the predictive value of said technique as a clinical screening tool was not analysed. Furthermore, genetic testing of serum and bodily fluids such as urine and saliva would reduce the need for biopsies to detect cancer and would thus be the most effective screening or monitoring tool. However the development of such tests requires an extremely high degree of sensitivity

and specificity. Analysis of GSTP1 gene hypermethylation (Cairns P, Esteller M, Herman JG, Schoenberg M, Jeronimo C, Sanchez-Cespedes M, et al. Molecular detection of prostate cancer in urine by GSTP1 hypermethylation. Clin Cancer Res 2001;7:2727-30.) in urine sediment of prostate cancer patients showed that only 6 out of 22 individuals with elevated methylation levels in biopsied tumors showed corresponding hypermethylation in urine samples.

Multifactorial approach. Cancer diagnostics has traditionally relied upon the detection of single molecular markers (e.g. gene mutations, elevated PSA levels). Unfortunately, cancer is a disease state in which single markers have typically failed to detect or differentiate many forms of the disease. Thus, assays that recognize only a single marker have been shown to be of limited predictive value. A fundamental aspect of this invention is that methylation based cancer diagnostics and the screening, diagnosis, and therapeutic monitoring of such diseases will provide significant improvements over the state-of-the-art that uses single marker analyses by the use of a selection of multiple markers. The multiplexed analytical approach is particularly well suited for cancer diagnostics since cancer is not a simple disease, this multi-factorial "panel" approach is consistent with the heterogeneous nature of cancer, both cytologically and clinically.

Key to the successful implementation of a panel approach to methylation based diagnostic tests is the design and development of optimized panels of markers that can characterize and distinguish disease states. This patent application describes an efficient and unique panel of genes the methylation analysis of one or a combination of the members of the panel enabling the detection of cell proliferative disorders of the prostate with a particularly high sensitivity, specificity and/or predictive value.

Pronounced need in the art. Therefore, in view of the incidence of prostate hyperplasia (50% of men aged 50 or above, and 95% of men aged 75 or above) and prostate cancer (180 per 100,000), there is a substantial need in the art for the development of molecular markers that could be used to

effectively distinguish among benign hyperplasia of the prostate and different grades of prostate cancer. Additionally, there is a pronounced need in the art for the development of molecular markers that could be used to provide sensitive, accurate and non-invasive methods (as opposed to, e.g., biopsy and transrectal ultrasound) for the diagnosis, prognosis and treatment of prostate cell proliferative disorders.

SUMMARY OF THE INVENTION

The disclosed invention provides a means for detection of or differentiation between prostate cell proliferative disorders by analysis of a gene panel, with a sensitivity and specificity suitable for use in a body fluid or serum assay. The present invention provides novel methods for detecting or distinguishing between prostate cell proliferative disorders with a sensitivity of greater than 30% and a specificity of greater than 65%. Said method is most preferably utilised for detecting or detecting and distinguishing between prostate cell proliferative disorders. The invention provides a method for the analysis of biological samples for features associated with the development of prostate cell proliferative disorders, the method characterised in that at least one nucleic acid, or a fragment thereof, from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 30 is/are contacted with a reagent or series of reagents capable of distinguishing between methylated and non methylated CpG dinucleotides within the genomic sequence, or sequences of interest.

The present invention provides a method for ascertaining genetic and/or epigenetic parameters of genomic DNA. The method has utility for the improved diagnosis, treatment and monitoring of prostate cell proliferative disorders, more specifically by enabling the improved identification of and differentiation between subclasses of said disorder and the genetic predisposition to said disorders. The invention presents improvements over the state of the art in that it enables a more specific and sensitive classification of prostate cell proliferative disorders than that achieved by

currently used tests thereby allowing for improved and informed treatment of patients.

Preferably, the source of the test sample is selected from the group consisting of cells or cell lines, histological slides, biopsies, paraffin-embedded tissue, bodily fluids, ejaculate, urine, blood, and combinations thereof. Preferably, the source is biopsies, bodily fluids, ejaculate, urine, or blood.

Specifically, the present invention provides a method for detecting prostate cell proliferative disorders with a sensitivity of greater than 30% and a specificity of greater than 65%, comprising: obtaining a biological sample comprising genomic nucleic acid(s); contacting the nucleic acid(s), or a fragment thereof, with one reagent or a plurality of reagents sufficient for distinguishing between methylated and non methylated CpG dinucleotide sequences within a target sequence of the subject nucleic acid, wherein the target sequence comprises, or hybridizes under stringent conditions to, a sequence comprising at least 16 contiguous nucleotides of SEQ ID NO: 1 to 30, said contiguous nucleotides comprising at least one CpG dinucleotide sequence; and determining, based at least in part on said distinguishing, the methylation state of at least one target CpG dinucleotide sequence, or an average, or a value reflecting an average methylation state of a plurality of target CpG dinucleotide sequences. Preferably, distinguishing between methylated and non methylated CpG dinucleotide sequences within the target sequence comprises methylation state-dependent conversion or non-conversion of at least one such CpG dinucleotide sequence to the corresponding converted or non-converted dinucleotide sequence within a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and contiguous regions thereof corresponding to the target sequence.

Additional embodiments provide a method for the detection of prostate cell proliferative disorders with a sensitivity of greater than 30% and a specificity of greater than 65%, comprising: obtaining a biological sample having subject genomic DNA; extracting the genomic DNA; treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms

of hybridization properties; contacting the treated genomic DNA, or the treated fragment thereof, with an amplification enzyme and at least two primers comprising, in each case a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, wherein the treated DNA or the fragment thereof is either amplified to produce an amplificate, or is not amplified; and determining, based on a presence or absence of, or on a property of said amplificate, the methylation state of at least one CpG dinucleotide sequence selected from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30, or an average, or a value reflecting an average methylation state of a plurality of CpG dinucleotide sequences thereof. Preferably, at least one such hybridizing nucleic acid molecule or peptide nucleic acid molecule is bound to a solid phase. Preferably, determining comprises use of at least two methods selected from the group consisting of: hybridizing at least one nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof; hybridizing at least one nucleic acid molecule, bound to a solid phase, comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof; hybridizing at least one nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, and extending at least one such hybridized nucleic acid molecule by at least one nucleotide base; and sequencing of the amplificate.

Additional embodiments provide novel genomic and chemically modified nucleic acid sequences, as well as oligonucleotides and/or PNA-oligomers for analysis of cytosine methylation

patterns within sequences from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30.

BRIEF DESCRIPTION OF THE DRAWINGS

DETAILED DESCRIPTION OF THE INVENTION

Definitions:

The term "Observed/Expected Ratio" ("O/E Ratio") refers to the frequency of CpG dinucleotides within a particular DNA sequence, and corresponds to the $\frac{\text{number of CpG sites}}{\text{number of C bases} \times \text{number of G bases}} \times \text{band length}$ for each fragment.

The term "CpG island" refers to a contiguous region of genomic DNA that satisfies the criteria of (1) having a frequency of CpG dinucleotides corresponding to an "Observed/Expected Ratio" >0.6 , and (2) having a "GC Content" >0.5 . CpG islands are typically, but not always, between about 0.2 to about 1 kb in length.

The term "methylation state" or "methylation status" refers to the presence or absence of 5-methylcytosine ("5-mCyt") at one or a plurality of CpG dinucleotides within a DNA sequence. Methylation states at one or more particular palindromic CpG methylation sites (each having two CpG CpG dinucleotide sequences) within a DNA sequence include "unmethylated," "fully-methylated" and "hemi-methylated."

The term "hemi-methylation" or "hemimethylation" refers to the methylation state of a palindromic CpG methylation site, where only a single cytosine in one of the two CpG dinucleotide sequences of the palindromic CpG methylation site is methylated (e.g., 5'-CC^MGG-3' (top strand): 3'-GGCC-5' (bottom strand)).

The term "hypermethylation" refers to the average methylation state corresponding to an *increased* presence of 5-mCyt at one or a plurality of CpG dinucleotides within a DNA sequence of a test DNA sample, relative to the amount of 5-mCyt found at corresponding CpG dinucleotides within a normal control DNA sample.

The term "hypomethylation" refers to the average methylation state corresponding to a *decreased* presence of 5-mCyt at one or a plurality of CpG dinucleotides within a DNA sequence of a

test DNA sample, relative to the amount of 5-mCyt found at corresponding CpG dinucleotides within a normal control DNA sample.

The term "microarray" refers broadly to both "DNA microarrays," and "DNA chip(s)," as recognized in the art, encompasses all art-recognized solid supports, and encompasses all methods for affixing nucleic acid molecules thereto or synthesis of nucleic acids thereon.

"Genetic parameters" are mutations and polymorphisms of genes and sequences further required for their regulation. To be designated as mutations are, in particular, insertions, deletions, point mutations, inversions and polymorphisms and, particularly preferred, SNPs (single nucleotide polymorphisms).

"Epigenetic parameters" are, in particular, cytosine methylations. Further epigenetic parameters include, for example, the acetylation of histones which, however, cannot be directly analyzed using the described method but which, in turn, correlate with the DNA methylation.

The term "bisulfite reagent" refers to a reagent comprising bisulfite, disulfite, hydrogen sulfite or combinations thereof, useful as disclosed herein to distinguish between methylated and unmethylated CpG dinucleotide sequences.

The term "Methylation assay" refers to any assay for determining the methylation state of one or more CpG dinucleotide sequences within a sequence of DNA.

The term "MS.AP-PCR" (Methylation-Sensitive Arbitrarily-Primed Polymerase Chain Reaction) refers to the art-recognized technology that allows for a global scan of the genome using CG-rich primers to focus on the regions most likely to contain CpG dinucleotides, and described by Gonzalgo et al., *Cancer Research* 57:594-599, 1997.

The term "MethyLight™" refers to the art-recognized fluorescence-based real-time PCR technique described by Eads et al., *Cancer Res.* 59:2302-2306, 1999.

The term "HeavyMethyl™" assay, in the embodiment thereof implemented herein, refers to a HeavyMethyl™ refer to the use of methylation specific *blocking* probes covering CpG positions between the amplification primers.

The term "MS-SNuPE" (Methylation-sensitive Single Nucleotide Primer Extension) refers to the art-recognized assay described by Gonzalgo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997.

The term "MSP" (Methylation-specific PCR) refers to the art-recognized methylation assay described by Herman et al. *Proc. Natl. Acad. Sci. USA* 93:9821-9826, 1996, and by US Patent No. 5,786,146.

The term "COBRA" (Combined Bisulfite Restriction Analysis) refers to the art-recognized methylation assay described by Xiong & Laird, *Nucleic Acids Res.* 25:2532-2534, 1997.

The term "MCA" (Methylated CpG Island Amplification) refers to the methylation assay described by Toyota et al., *Cancer Res.* 59:2307-12, 1999, and in WO 00/26401A1.

The term "hybridization" is to be understood as a bond of an oligonucleotide to a complementary sequence along the lines of the Watson-Crick base pairings in the sample DNA, forming a duplex structure.

"Stringent hybridization conditions," as defined herein, involve hybridizing at 68°C in 5x SSC/5x Denhardt's solution/1.0% SDS, and washing in 0.2x SSC/0.1% SDS at room temperature, or involve the art-recognized equivalent thereof (e.g., conditions in which a hybridization is carried out at 60°C in 2.5 x SSC buffer, followed by several washing steps at 37°C in a low buffer concentration, and remains stable). Moderately stringent conditions, as defined herein, involve including washing in 3x SSC at 42°C, or the art-recognized equivalent thereof. The parameters of salt concentration and temperature can be varied to achieve the optimal level of identity between the probe and the target nucleic acid. Guidance regarding such conditions is available in the art, for example, by Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, N.Y.; and Ausubel et al. (eds.), 1995, *Current Protocols in Molecular Biology*, (John Wiley & Sons, N.Y.) at Unit 2.10.

The terms 'sensitivity' and 'specificity' refer to values calculated with reference to a sample set of male patients with an average age of 65 and a mixed ethnic range including caucasian and african american.

Overview:

The present invention provides for molecular genetic markers that have novel utility for the analysis of methylation patterns associated with the development of prostate cell proliferative disorders with a sensitivity of greater than 30% and a specificity of greater than 65%. Said markers may be used for detecting or distinguishing between prostate cell proliferative disorders, thereby providing improved means for the classification and treatment of said disorders. The markers according to the present invention are analysed in the form of a 'panel' wherein the methylation of one or more genetic sequences of the genes

Bisulfite modification of DNA is an art-recognized tool used to assess CpG methylation status. 5-methylcytosine is the most frequent covalent base modification in the DNA of eukaryotic cells. It plays a role, for example, in the regulation of the transcription, in genetic imprinting, and in tumorigenesis. Therefore, the identification of 5-methylcytosine as a component of genetic information is of considerable interest. However, 5-methylcytosine positions cannot be identified by sequencing, because 5-methylcytosine has the same base pairing behavior as cytosine. Moreover, the epigenetic information carried by 5-methylcytosine is completely lost during, e.g., PCR amplification.

The most frequently used method for analyzing DNA for the presence of 5-methylcytosine is based upon the specific reaction of bisulfite with cytosine whereby, upon subsequent alkaline hydrolysis, cytosine is converted to uracil which corresponds to thymine in its base pairing behavior. Significantly, however, 5-methylcytosine remains unmodified under these conditions. Consequently, the original DNA is converted in such a manner that methylcytosine, which originally could not be distinguished from cytosine by its hybridization behavior, can now be detected as the only remaining cytosine using standard, art-recognized molecular biological techniques, for example, by amplification and hybridization, or by sequencing. All of these techniques are based on differential base pairing properties, which can now be fully exploited.

The prior art, in terms of sensitivity, is defined by a method comprising enclosing the DNA to be analyzed in an agarose matrix, thereby preventing the diffusion and renaturation of the DNA (bisulfite only reacts with single-stranded DNA), and replacing all precipitation and purification steps with fast dialysis (Olek A, et al., A modified and improved method for bisulfite based cytosine methylation analysis, *Nucleic Acids Res.* 24:5064-6, 1996). It is thus possible to analyze individual cells for methylation status, illustrating the utility and sensitivity of the method. An overview of art-recognized methods for detecting 5-methylcytosine is provided by Rein, T., et al., *Nucleic Acids Res.*, 26:2255, 1998.

The bisulfite technique, barring few exceptions (e.g., Zeschnigk M, et al., *Eur J Hum Genet.* 5:94-98, 1997), is currently only used in research. In all instances, short, specific fragments of a known gene are amplified subsequent to a bisulfite treatment, and either completely sequenced (Olek & Walter, *Nat Genet.* 1997 17:275-6, 1997), subjected to one or more primer extension reactions (Gonzalzo & Jones, *Nucleic Acids Res.*, 25:2529-31, 1997; WO 95/00669; U.S. Patent No. 6,251,594) to analyze individual cytosine positions, or treated by enzymatic digestion (Xiong & Laird, *Nucleic Acids Res.*, 25:2532-4, 1997). Detection by hybridization has also been described in the art (Olek et al., WO 99/28498). Additionally, use of the bisulfite technique for methylation detection with respect to individual genes has been described (Grigg & Clark, *Bioessays*, 16:431-6, 1994; Zeschnigk M, et al., *Hum Mol Genet.*, 6:387-95, 1997; Feil R, et al., *Nucleic Acids Res.*, 22:695-, 1994; Martin V, et al., *Gene*, 157:261-4, 1995; WO 9746705 and WO 9515373).

The present invention provides for the use of the bisulfite technique, in combination with one or more methylation assays, for determination of the methylation status of CpG dinucleotide sequences within sequences from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30. According to the present invention, determination of the methylation status of CpG dinucleotide sequences within sequences from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 has diagnostic and prognostic utility.

Methylation Assay Procedures. Various methylation assay procedures are known in the art, and can be used in conjunction with the present invention. These assays allow for determination of the methylation state of one or a plurality of CpG dinucleotides (e.g., CpG islands) within a DNA sequence. Such assays involve, among other techniques, DNA sequencing of bisulfite-treated DNA, PCR (for sequence-specific amplification), Southern blot analysis, and use of methylation-sensitive restriction enzymes.

For example, genomic sequencing has been simplified for analysis of DNA methylation patterns and 5-methylcytosine distribution by using bisulfite treatment (Frommer et al., *Proc. Natl. Acad. Sci. USA* 89:1827-1831, 1992). Additionally, restriction enzyme digestion of PCR products amplified from bisulfite-converted DNA is used, e.g., the method described by Sadri & Hornsby (*Nucl. Acids Res.* 24:5058-5059, 1996), or COBRA (Combined Bisulfite Restriction Analysis) (Xiong & Laird, *Nucleic Acids Res.* 25:2532-2534, 1997).

COBRA. COBRA analysis is a quantitative methylation assay useful for determining DNA methylation levels at specific gene loci in small amounts of genomic DNA (Xiong & Laird, *Nucleic Acids Res.* 25:2532-2534, 1997). Briefly, restriction enzyme digestion is used to reveal methylation-dependent sequence differences in PCR products of sodium bisulfite-treated DNA. Methylation-dependent sequence differences are first introduced into the genomic DNA by standard bisulfite treatment according to the procedure described by Frommer et al. (*Proc. Natl. Acad. Sci. USA* 89:1827-1831, 1992). PCR amplification of the bisulfite converted DNA is then performed using primers specific for the interested CpG islands, followed by restriction endonuclease digestion, gel electrophoresis, and detection using specific, labeled hybridization probes. Methylation levels in the original DNA sample are represented by the relative amounts of digested and undigested PCR product in a linearly quantitative fashion across a wide spectrum of DNA methylation levels. In addition, this technique can be reliably applied to DNA obtained from microdissected paraffin-embedded tissue samples. Typical reagents (e.g., as might be found in a typical COBRA-based kit) for COBRA analysis may include, but are not limited to: PCR primers for specific gene

(or methylation-altered DNA sequence or CpG island); restriction enzyme and appropriate buffer; gene-hybridization oligo; control hybridization oligo; kinase labeling kit for oligo probe; and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kits (e.g., precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

Preferably, assays such as "MethyLight•" (a fluorescence-based real-time PCR technique) (Eads et al., *Cancer Res.* 59:2302-2306, 1999), Ms-SNuPE (Methylation-sensitive Single Nucleotide Primer Extension) reactions (Gonzalzo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997), methylation-specific PCR ("MSP"; Herman et al., *Proc. Natl. Acad. Sci. USA* 93:9821-9826, 1996; US Patent No. 5,786,146), and methylated CpG island amplification ("MCA"; Toyota et al., *Cancer Res.* 59:2307-12, 1999) are used alone or in combination with other of these methods.

MethyLight•. The MethyLight• assay is a high-throughput quantitative methylation assay that utilizes fluorescence-based real-time PCR (TaqMan••) technology that requires no further manipulations after the PCR step (Eads et al., *Cancer Res.* 59:2302-2306, 1999). Briefly, the MethyLight• process begins with a mixed sample of genomic DNA that is converted, in a sodium bisulfite reaction, to a mixed pool of methylation-dependent sequence differences according to standard procedures (the bisulfite process converts unmethylated cytosine residues to uracil). Fluorescence-based PCR is then performed either in an "unbiased" (with primers that do not overlap known CpG methylation sites) PCR reaction, or in a "biased" (with PCR primers that overlap known CpG dinucleotides) reaction. Sequence discrimination can occur either at the level of the amplification process or at the level of the fluorescence detection process, or both.

The MethyLight• assay may be used as a quantitative test for methylation patterns in the genomic DNA sample, wherein sequence discrimination occurs at the level of probe hybridization. In this quantitative version, the PCR reaction provides for unbiased amplification in the presence of a fluorescent probe that overlaps a particular putative

methylation site. An unbiased control for the amount of input DNA is provided by a reaction in which neither the primers, nor the probe overlies any CpG dinucleotides. Alternatively, a qualitative test for genomic methylation is achieved by probing of the biased PCR pool with either control oligonucleotides that do not "cover" known methylation sites (a fluorescence-based version of the "MSP" technique), or with oligonucleotides covering potential methylation sites.

The MethyLight[®] process can be used with a "TaqMan[®]" probe in the amplification process. For example, double-stranded genomic DNA is treated with sodium bisulfite and subjected to one of two sets of PCR reactions using TaqMan[®] probes; e.g., with either biased primers and TaqMan[®] probe, or unbiased primers and TaqMan[®] probe. The TaqMan[®] probe is dual-labeled with fluorescent "reporter" and "quencher" molecules, and is designed to be specific for a relatively high GC content region so that it melts out at about 10°C higher temperature in the PCR cycle than the forward or reverse primers. This allows the TaqMan[®] probe to remain fully hybridized during the PCR annealing/extension step. As the Taq polymerase enzymatically synthesizes a new strand during PCR, it will eventually reach the annealed TaqMan[®] probe. The Taq polymerase 5' to 3' endonuclease activity will then displace the TaqMan[®] probe by digesting it to release the fluorescent reporter molecule for quantitative detection of its now unquenched signal using a real-time fluorescent detection system.

Typical reagents (e.g., as might be found in a typical MethyLight[®]-based kit) for MethyLight[®] analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); TaqMan[®] probes; optimized PCR buffers and deoxynucleotides; and Taq polymerase.

Ms-SNuPE. The Ms-SNuPE technique is a quantitative method for assessing methylation differences at specific CpG sites based on bisulfite treatment of DNA, followed by single-nucleotide primer extension (Gonzalzo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997). Briefly, genomic DNA is reacted with sodium bisulfite to convert unmethylated cytosine to uracil while leaving 5-methylcytosine unchanged. Amplification of the desired target sequence is then performed using PCR primers

specific for bisulfite-converted DNA, and the resulting product is isolated and used as a template for methylation analysis at the CpG site(s) of interest. Small amounts of DNA can be analyzed (e.g., microdissected pathology sections), and it avoids utilization of restriction enzymes for determining the methylation status at CpG sites.

Typical reagents (e.g., as might be found in a typical Ms-SNuPE-based kit) for Ms-SNuPE analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); optimized PCR buffers and deoxynucleotides; gel extraction kit; positive control primers; Ms-SNuPE primers for specific gene; reaction buffer (for the Ms-SNuPE reaction); and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kit (e.g., precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

MSP. MSP (methylation-specific PCR) allows for assessing the methylation status of virtually any group of CpG sites within a CpG island, independent of the use of methylation-sensitive restriction enzymes (Herman et al. *Proc. Natl. Acad. Sci. USA* 93:9821-9826, 1996; US Patent No. 5,786,146). Briefly, DNA is modified by sodium bisulfite converting all unmethylated, but not methylated cytosines to uracil, and subsequently amplified with primers specific for methylated versus unmethylated DNA. MSP requires only small quantities of DNA, is sensitive to 0.1% methylated alleles of a given CpG island locus, and can be performed on DNA extracted from paraffin-embedded samples. Typical reagents (e.g., as might be found in a typical MSP-based kit) for MSP analysis may include, but are not limited to: methylated and unmethylated PCR primers for specific gene (or methylation-altered DNA sequence or CpG island), optimized PCR buffers and deoxynucleotides, and specific probes.

MCA. The MCA technique is a method that can be used to screen for altered methylation patterns in genomic DNA, and to isolate specific sequences associated with these changes (Toyota et al., *Cancer Res.* 59:2307-12, 1999). Briefly, restriction enzymes with different sensitivities to cytosine methylation in their recognition sites are used to digest

genomic DNAs from primary tumors, cell lines, and normal tissues prior to arbitrarily primed PCR amplification. Fragments that show differential methylation are cloned and sequenced after resolving the PCR products on high-resolution polyacrylamide gels. The cloned fragments are then used as probes for Southern analysis to confirm differential methylation of these regions. Typical reagents (e.g., as might be found in a typical MCA-based kit) for MCA analysis may include, but are not limited to: PCR primers for arbitrary priming Genomic DNA; PCR buffers and nucleotides, restriction enzymes and appropriate buffers; gene-hybridization oligos or probes; control hybridization oligos or probes.

HeavyMethyl. The HeavyMethyl techniques is a means for selectively amplifying methylated as opposed to non-methylated DNA (or vice versa). Blocker oligonucleotides specific to either methylated or unmethylated versions of a bisulfite treated target sequence are hybridised to the treated nucleic acids. The sample is then enzymatically amplified, wherein the hybridisation of the blocker oligonucleotides hinders amplification of the nucleic acid strand to which it is bound. Typical reagents (e.g., as might be found in a typical HeavyMethyl-based kit) for HeavyMethyl analysis may include, but are not limited to: methylated or unmethylated blocker oligonucleotides for specific gene (or methylation-altered DNA sequence or CpG island), optimized PCR buffers and deoxynucleotides, and specific probes and primers.

GENOMIC SEQUENCES ACCORDING TO SEQ ID NO: 1 TO SEQ ID NO: 4 AND SEQ ID NOS: 29 & 30, AND TREATED VARIANTS THEREOF ACCORDING TO SEQ ID NO: 5 TO SEQ ID NO: 28, WERE DETERMINED TO HAVE UTILITY FOR DETECTING OR DISTINGUISHING BETWEEN OR AMONG PROSTATE CELL PROLIFERATIVE DISORDERS. .

The present invention is based upon the analysis of methylation levels within one or more genes taken from the group consisting GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions and sequences thereof according to Table 5.

Particular embodiments of the present invention provide a novel application of the analysis of methylation levels and/or patterns within said genes and/or sequences that enables a precise detection, characterisation and/or treatment of prostate cell proliferative disorders. Early detection of prostate cell proliferative disorders is directly linked with disease prognosis, and the disclosed method thereby enables the physician and patient to make better and more informed treatment decisions. The methods disclosed according to the invention enable the detection and characterisation of prostate cell proliferative disorders with improved sensitivity and/or specificity of with a sensitivity of greater than 30% and a specificity of greater than 65%.

FURTHER IMPROVEMENTS

The present invention provides novel uses for genomic sequences selected from the group consisting of SEQ ID NO: 1 TO SEQ ID NO: 4 AND SEQ ID NOS: 29 & 30. Additional embodiments provide modified variants of SEQ ID NO: 1 TO SEQ ID NO: 4 AND SEQ ID NOS: 29 & 30, as well as oligonucleotides and/or PNA-oligomers for analysis of cytosine methylation patterns within SEQ ID NO: 1 TO SEQ ID NO: 4 AND SEQ ID NOS: 29 & 30.

An objective of the invention comprises analysis of the methylation state of one or more CpG dinucleotides within at least one of the genomic sequences selected from the group consisting of SEQ ID NO: 1 TO SEQ ID NO: 4 AND SEQ ID NOS: 29 & 30 and sequences complementary thereto.

The disclosed invention provides treated nucleic acids, derived from genomic SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30, wherein the treatment is suitable to convert at least one unmethylated cytosine base of the genomic DNA sequence to uracil or another base that is detectably dissimilar to cytosine in terms of hybridization. The genomic sequences in question may comprise one, or more, consecutive or random methylated CpG positions. Said treatment preferably comprises use of a reagent selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof. In a preferred embodiment of the invention, the objective comprises analysis of a modified nucleic acid comprising a sequence of at least 16 contiguous nucleotide bases in length of a sequence selected from the group

consisting of SEQ ID NO: 5 TO SEQ ID NO: 28, wherein said sequence comprises at least one CpG, TpA or CpA dinucleotide and sequences complementary thereto. The sequences of SEQ ID NO: 5 TO SEQ ID NO: 28 provide modified versions of the nucleic acid according to SEQ ID NO: 1 TO SEQ ID NO: 4 AND SEQ ID NOS: 29 & 30, wherein the modification of each genomic sequence results in the synthesis of a nucleic acid having a sequence that is unique and distinct from said genomic sequence as follows. For each sense strand genomic DNA, e.g., SEQ ID NO:1, four converted versions are disclosed. A first version wherein "C".."T," but "CpG" remains "CpG" (i.e., corresponds to case where, for the genomic sequence, all "C" residues of CpG dinucleotide sequences are methylated and are thus not converted); a second version discloses the complement of the disclosed genomic DNA sequence (i.e. antisense strand), wherein "C".."T," but "CpG" remains "CpG" (i.e., corresponds to case where, for all "C" residues of CpG dinucleotide sequences are methylated and are thus not converted). The 'upmethylated' converted sequences of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 correspond to SEQ ID NO: 5 to SEQ ID NO: 12 and SEQ ID NO: 21 to 24. A third chemically converted version of each genomic sequences is provided, wherein "C".."T" for all "C" residues, including those of "CpG" dinucleotide sequences (i.e., corresponds to case where, for the genomic sequences, all "C" residues of CpG dinucleotide sequences are unmethylated); a final chemically converted version of each sequence, discloses the complement of the disclosed genomic DNA sequence (i.e. antisense strand), wherein "C".."T" for all "C" residues, including those of "CpG" dinucleotide sequences (i.e., corresponds to case where, for the complement (antisense strand) of each genomic sequence, all "C" residues of CpG dinucleotide sequences are unmethylated). The 'downmethylated' converted sequences of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 correspond to SEQ ID NO: 13 to SEQ ID NO: 20 and SEQ ID NO: 25 to 28.

In an alternative preferred embodiment, such analysis comprises the use of an oligonucleotide or oligomer for detecting the cytosine methylation state within genomic or pretreated (chemically modified) DNA, according to SEQ ID NO: 1

to SEQ ID NO: 30. Said oligonucleotide or oligomer comprising a nucleic acid sequence having a length of at least nine (9) nucleotides which hybridizes, under moderately stringent or stringent conditions (as defined herein above), to a pretreated nucleic acid sequence according to SEQ ID NO: 5 to SEQ ID NO: 28 and/or sequences complementary thereto, or to a genomic sequence according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 and/or sequences complementary thereto.

Thus, the present invention includes nucleic acid molecules (e.g., oligonucleotides and peptide nucleic acid (PNA) molecules (PNA-oligomers)) that hybridize under moderately stringent and/or stringent hybridization conditions to all or a portion of the sequences SEQ ID NO: 1 to SEQ ID NO: 30, or to the complements thereof. The hybridizing portion of the hybridizing nucleic acids is typically at least 9, 15, 20, 25, 30 or 35 nucleotides in length. However, longer molecules have inventive utility, and are thus within the scope of the present invention.

Preferably, the hybridizing portion of the inventive hybridizing nucleic acids is at least 95%, or at least 98%, or 100% identical to the sequence, or to a portion thereof of SEQ ID NO: 1 to SEQ ID NO: 30, or to the complements thereof.

Hybridizing nucleic acids of the type described herein can be used, for example, as a primer (e.g., a PCR primer), or a diagnostic and/or prognostic probe or primer. Preferably, hybridization of the oligonucleotide probe to a nucleic acid sample is performed under stringent conditions and the probe is 100% identical to the target sequence. Nucleic acid duplex or hybrid stability is expressed as the melting temperature or T_m , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions.

For target sequences that are related and substantially identical to the corresponding sequence of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 (such as allelic variants and SNPs), rather than identical, it is useful to first establish the lowest temperature at which only homologous hybridization occurs with a particular concentration of salt (e.g., SSC or SSPE). Then, assuming that 1% mismatching results in a 1°C

decrease in the T_m , the temperature of the final wash in the hybridization reaction is reduced accordingly (for example, if sequences having > 95% identity with the probe are sought, the final wash temperature is decreased by 5°C). In practice, the change in T_m can be between 0.5°C and 1.5°C per 1% mismatch.

Examples of inventive oligonucleotides of length X (in nucleotides), as indicated by polynucleotide positions with reference to, e.g., SEQ ID NO:1, include those corresponding to sets (sense and antisense sets) of consecutively overlapping oligonucleotides of length X, where the oligonucleotides within each consecutively overlapping set (corresponding to a given X value) are defined as the finite set of Z oligonucleotides from nucleotide positions:

n to (n + (X-1));

where n=1, 2, 3,...(Y-(X-1));

where Y equals the length (nucleotides or base pairs) of SEQ ID NO:1 (VALUE TO REFLECT LENGTH OF SEQ ID NO:1);

where X equals the common length (in nucleotides) of each oligonucleotide in the set (e.g., X=20 for a set of consecutively overlapping 20-mers); and

where the number (Z) of consecutively overlapping oligomers of length X for a given SEQ ID NO of length Y is equal to Y-(X-1). For example Z= VALUE TO REFLECT LENGTH OF SEQ ID NO:1 -19= VALUE TO REFLECT LENGTH OF SEQ ID NO:1 for either sense or antisense sets of SEQ ID NO:1, where X=20.

Preferably, the set is limited to those oligomers that comprise at least one CpG, TpG or CpA dinucleotide.

Examples of inventive 20-mer oligonucleotides include the following set of VALUE TO REFLECT LENGTH OF SEQ ID NO:1 oligomers (and the antisense set complementary thereto), indicated by polynucleotide positions with reference to SEQ ID NO:1 1-20, 2-21, 3-22, 4-23, 5-24,

Preferably, the set is limited to those oligomers that comprise at least one CpG, TpG or CpA dinucleotide.

The present invention encompasses, for each of SEQ ID NO: 1 to SEQ ID NO: 30 (sense and antisense), multiple consecutively overlapping sets of oligonucleotides or modified

oligonucleotides of length X, where, e.g., X= 9, 10, 17, 20, 22, 23, 25, 27, 30 or 35 nucleotides.

The oligonucleotides or oligomers according to the present invention constitute effective tools useful to ascertain genetic and epigenetic parameters of the genomic sequence corresponding to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30. Preferred sets of such oligonucleotides or modified oligonucleotides of length X are those consecutively overlapping sets of oligomers corresponding to SEQ ID NO: 1 to SEQ ID NO: 30 (and to the complements thereof). Preferably, said oligomers comprise at least one CpG, TpG or CpA dinucleotide.

Particularly preferred oligonucleotides or oligomers according to the present invention are those in which the cytosine of the CpG dinucleotide (or of the corresponding converted TpG or CpA dinucleotide) sequences is within the middle third of the oligonucleotide; that is, where the oligonucleotide is, for example, 13 bases in length, the CpG, TpG or CpA dinucleotide is positioned within the fifth to ninth nucleotide from the 5'-end.

The oligonucleotides of the invention can also be modified by chemically linking the oligonucleotide to one or more moieties or conjugates to enhance the activity, stability or detection of the oligonucleotide. Such moieties or conjugates include chromophores, fluorophors, lipids such as cholesterol, cholic acid, thioether, aliphatic chains, phospholipids, polyamines, polyethylene glycol (PEG), palmityl moieties, and others as disclosed in, for example, United States Patent Numbers 5,514,758, 5,565,552, 5,567,810, 5,574,142, 5,585,481, 5,587,371, 5,597,696 and 5,958,773. The probes may also exist in the form of a PNA (peptide nucleic acid) which has particularly preferred pairing properties. Thus, the oligonucleotide may include other appended groups such as peptides, and may include hybridization-triggered cleavage agents (Krol et al., *BioTechniques* 6:958-976, 1988) or intercalating agents (Zon, *Pharm. Res.* 5:539-549, 1988). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a chromophore, fluorophor, peptide, hybridization-triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The oligonucleotide may also comprise at least one art-recognized modified sugar and/or base moiety, or may comprise a modified backbone or non-natural internucleoside linkage.

The oligonucleotides or oligomers according to particular embodiments of the present invention are typically used in 'sets,' which contain at least one oligomer for analysis of each of the CpG dinucleotides of genomic sequence SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 and sequences complementary thereto, or to the corresponding CpG, TpG or CpA dinucleotide within a sequence of the pretreated nucleic acids according to SEQ ID NO: 5 to SEQ ID NO: 28 and sequences complementary thereto. However, it is anticipated that for economic or other factors it may be preferable to analyze a limited selection of the CpG dinucleotides within said sequences, and the content of the set of oligonucleotides is altered accordingly.

Therefore, in particular embodiments, the present invention provides a set of at least two (2) (oligonucleotides and/or PNA-oligomers) useful for detecting the cytosine methylation state in pretreated genomic DNA (SEQ ID NO: 5 to SEQ ID NO: 28), or in genomic DNA (SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 and sequences complementary thereto). These probes enable diagnosis, classification and/or therapy of genetic and epigenetic parameters of prostate cell proliferative disorders. The set of oligomers may also be used for detecting single nucleotide polymorphisms (SNPs) in pretreated genomic DNA (SEQ ID NO: 5 to SEQ ID NO: 28), or in genomic DNA (SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 and sequences complementary thereto).

In preferred embodiments, at least one, and more preferably all members of a set of oligonucleotides is bound to a solid phase.

In further embodiments, the present invention provides a set of at least two (2) oligonucleotides that are used as 'primer' oligonucleotides for amplifying DNA sequences of one of SEQ ID NO: 1 to SEQ ID NO: 30 and sequences complementary thereto, or segments thereof.

It is anticipated that the oligonucleotides may constitute all or part of an "array" or "DNA chip" (i.e., an arrangement of different oligonucleotides and/or PNA-oligomers bound to a

solid phase). Such an array of different oligonucleotide- and/or PNA-oligomer sequences can be characterized, for example, in that it is arranged on the solid phase in the form of a rectangular or hexagonal lattice. The solid-phase surface may be composed of silicon, glass, polystyrene, aluminum, steel, iron, copper, nickel, silver, or gold. Nitrocellulose as well as plastics such as nylon, which can exist in the form of pellets or also as resin matrices, may also be used. An overview of the Prior Art in oligomer array manufacturing can be gathered from a special edition of *Nature Genetics* (*Nature Genetics Supplement*, Volume 21, January 1999, and from the literature cited therein). Fluorescently labeled probes are often used for the scanning of immobilized DNA arrays. The simple attachment of Cy3 and Cy5 dyes to the 5'-OH of the specific probe are particularly suitable for fluorescence labels. The detection of the fluorescence of the hybridized probes may be carried out, for example, via a confocal microscope. Cy3 and Cy5 dyes, besides many others, are commercially available.

It is particularly preferred that the oligomers according to the invention are utilised for at least one of: detection of; detection and differentiation between or among subclasses of; diagnosis of; prognosis of; treatment of; monitoring of; and treatment and monitoring of prostate cell proliferative disorders. This is enabled by use of said sets for the detection or detection and differentiation of prostate cell proliferative disorders.

The present invention further provides a method for ascertaining genetic and/or epigenetic parameters of the genes GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions including genomic sequences according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 within a subject by analyzing cytosine methylation and single nucleotide polymorphisms. Said method comprising contacting a nucleic acid comprising one or more of the genes GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions including genomic sequences according to

SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 in a biological sample obtained from said subject with at least one reagent or a series of reagents, wherein said reagent or series of reagents, distinguishes between methylated and non-methylated CpG dinucleotides within the target nucleic acid. Preferably, said method comprises the following steps: In the *first step*, a sample of the tissue to be analysed is obtained. The source may be any suitable source, such as cell lines, histological slides, biopsies, tissue embedded in paraffin, bodily fluids, ejaculate, urine, blood and all possible combinations thereof. The DNA is then isolated from the sample. Extraction may be by means that are standard to one skilled in the art, including the use of commercially available kits, detergent lysates, sonification and vortexing with glass beads. Once the nucleic acids have been extracted, the genomic double stranded DNA is used in the analysis.

In the *second step* of the method, the genomic DNA sample is treated in such a manner that cytosine bases which are unmethylated at the 5'-position are converted to uracil, thymine, or another base which is dissimilar to cytosine in terms of hybridization behavior. This will be understood as 'pretreatment' herein.

The above described treatment of genomic DNA is preferably carried out with bisulfite (hydrogen sulfite, disulfite) and subsequent alkaline hydrolysis which results in a conversion of non-methylated cytosine nucleobases to uracil or to another base which is dissimilar to cytosine in terms of base pairing behavior.

In the *third step* of the method, fragments of the pretreated DNA are amplified, using sets of primer oligonucleotides according to the present invention, and an amplification enzyme. The amplification of several DNA segments can be carried out simultaneously in one and the same reaction vessel. Typically, the amplification is carried out using a polymerase chain reaction (PCR). The set of primer oligonucleotides includes at least two oligonucleotides whose sequences are each reverse complementary, identical, or hybridize under stringent or highly stringent conditions to an at least 16-base-pair long segment of the base sequences of one

or more of SEQ ID NO: 5 to SEQ ID NO: 28 and sequences complementary thereto.

In an alternate embodiment of the method, the methylation status of preselected CpG positions within the nucleic acid sequences comprising one or more of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 may be detected by use of methylation-specific primer oligonucleotides. This technique (MSP) has been described in United States Patent No. 6,265,171 to Herman. The use of methylation status specific primers for the amplification of bisulfite treated DNA allows the differentiation between methylated and unmethylated nucleic acids. MSP primers pairs contain at least one primer which hybridizes to a bisulfite treated CpG dinucleotide. Therefore, the sequence of said primers comprises at least one CpG dinucleotide. MSP primers specific for non-methylated DNA contain a "T" at the 3' position of the C position in the CpG. Preferably, therefore, the base sequence of said primers is required to comprise a sequence having a length of at least 9 nucleotides which hybridizes to a pretreated nucleic acid sequence according to one of SEQ ID NO: 5 to SEQ ID NO: 28 and sequences complementary thereto, wherein the base sequence of said oligomers comprises at least one CpG dinucleotide.

A further preferred embodiment of the method comprises the use of blocker oligonucleotides. The use of such blocker oligonucleotides has been described by Yu et al., *Biotechniques* 23:714-720, 1997. Blocking probe oligonucleotides are hybridized to the bisulfite treated nucleic acid concurrently with the PCR primers. PCR amplification of the nucleic acid is terminated at the 5' position of the blocking probe, such that amplification of a nucleic acid is suppressed where the complementary sequence to the blocking probe is present. The probes may be designed to hybridize to the bisulfite treated nucleic acid in a methylation status specific manner. For example, for detection of methylated nucleic acids within a population of unmethylated nucleic acids, suppression of the amplification of nucleic acids which are unmethylated at the position in question would be carried out by the use of blocking probes comprising a 'CpA' or 'TpA' at the position in question, as opposed to a 'CpG' if the suppression of amplification of methylated nucleic acids is desired.

For PCR methods using blocker oligonucleotides, efficient disruption of polymerase-mediated amplification requires that blocker oligonucleotides not be elongated by the polymerase. Preferably, this is achieved through the use of blockers that are 3'-deoxyoligonucleotides, or oligonucleotides derivitized at the 3' position with other than a "free" hydroxyl group. For example, 3'-O-acetyl oligonucleotides are representative of a preferred class of blocker molecule.

Additionally, polymerase-mediated decomposition of the blocker oligonucleotides should be precluded. Preferably, such preclusion comprises either use of a polymerase lacking 5'-3' exonuclease activity, or use of modified blocker oligonucleotides having, for example, thioate bridges at the 5'-termini thereof that render the blocker molecule nuclease-resistant. Particular applications may not require such 5' modifications of the blocker. For example, if the blocker- and primer-binding sites overlap, thereby precluding binding of the primer (e.g., with excess blocker), degradation of the blocker oligonucleotide will be substantially precluded. This is because the polymerase will not extend the primer toward, and through (in the 5'-3' direction) the blocker—a process that normally results in degradation of the hybridized blocker oligonucleotide.

A particularly preferred blocker/PCR embodiment, for purposes of the present invention and as implemented herein, comprises the use of peptide nucleic acid (PNA) oligomers as blocking oligonucleotides. Such PNA blocker oligomers are ideally suited, because they are neither decomposed nor extended by the polymerase.

Preferably, therefore, the base sequence of said *blocking oligonucleotides* is required to comprise a sequence having a length of at least 9 nucleotides which hybridizes to a pretreated nucleic acid sequence according to one of SEQ ID NO: 5 to SEQ ID NO: 28 and sequences complementary thereto, wherein the base sequence of said oligonucleotides comprises at least one CpG, TpG or CpA dinucleotide.

The fragments obtained by means of the amplification can carry a directly or indirectly detectable label. Preferred are labels in the form of fluorescence labels, radionuclides, or

detachable molecule fragments having a typical mass which can be detected in a mass spectrometer. Where said labels are mass labels, it is preferred that the labeled amplificates have a single positive or negative net charge, allowing for better detectability in the mass spectrometer. The detection may be carried out and visualized by means of, e.g., matrix assisted laser desorption/ionization mass spectrometry (MALDI) or using electron spray mass spectrometry (ESI).

Matrix Assisted Laser Desorption/Ionization Mass Spectrometry (MALDI-TOF) is a very efficient development for the analysis of biomolecules (Karas & Hillenkamp, *Anal Chem.*, 60:2299-301, 1988). An analyte is embedded in a light-absorbing matrix. The matrix is evaporated by a short laser pulse thus transporting the analyte molecule into the vapour phase in an unfragmented manner. The analyte is ionized by collisions with matrix molecules. An applied voltage accelerates the ions into a field-free flight tube. Due to their different masses, the ions are accelerated at different rates. Smaller ions reach the detector sooner than bigger ones. MALDI-TOF spectrometry is well suited to the analysis of peptides and proteins. The analysis of nucleic acids is somewhat more difficult (Gut & Beck, *Current Innovations and Future Trends*, 1:147-57, 1995). The sensitivity with respect to nucleic acid analysis is approximately 100-times less than for peptides, and decreases disproportionally with increasing fragment size. Moreover, for nucleic acids having a multiply negatively charged backbone, the ionization process via the matrix is considerably less efficient. In MALDI-TOF spectrometry, the selection of the matrix plays an eminently important role. For desorption of peptides, several very efficient matrixes have been found which produce a very fine crystallisation. There are now several responsive matrixes for DNA, however, the difference in sensitivity between peptides and nucleic acids has not been reduced. This difference in sensitivity can be reduced, however, by chemically modifying the DNA in such a manner that it becomes more similar to a peptide. For example, phosphorothioate nucleic acids, in which the usual phosphates of the backbone are substituted with thiophosphates, can be converted into a charge-neutral DNA using simple alkylation chemistry (Gut & Beck, *Nucleic Acids*

Res. 23: 1367-73, 1995). The coupling of a charge tag to this modified DNA results in an increase in MALDI-TOF sensitivity to the same level as that found for peptides. A further advantage of charge tagging is the increased stability of the analysis against impurities, which makes the detection of unmodified substrates considerably more difficult.

In the *fourth step* of the method, the amplicates obtained during the third step of the method are analysed in order to ascertain the methylation status of the CpG dinucleotides prior to the treatment.

In embodiments where the amplicates were obtained by means of MSP amplification, the presence or absence of an amplicate is in itself indicative of the methylation state of the CpG positions covered by the primer, according to the base sequences of said primer.

Amplicates obtained by means of both standard and methylation specific PCR may be further analyzed by means of hybridization-based methods such as, but not limited to, array technology and probe based technologies as well as by means of techniques such as sequencing and template directed extension.

In one embodiment of the method, the amplicates synthesised in *step three* are subsequently hybridized to an array or a set of oligonucleotides and/or PNA probes. In this context, the hybridization takes place in the following manner: the set of probes used during the hybridization is preferably composed of at least 2 oligonucleotides or PNA-oligomers; in the process, the amplicates serve as probes which hybridize to oligonucleotides previously bonded to a solid phase; the non-hybridized fragments are subsequently removed; said oligonucleotides contain at least one base sequence having a length of at least 9 nucleotides which is reverse complementary or identical to a segment of the base sequences specified in the present Sequence Listing; and the segment comprises at least one CpG, TpG or CpA dinucleotide.

In a preferred embodiment, said dinucleotide is present in the central third of the oligomer. For example, wherein the oligomer comprises one CpG dinucleotide, said dinucleotide is preferably the fifth to ninth nucleotide from the 5'-end of a 13-mer. One oligonucleotide exists for the analysis of each CpG dinucleotide within the sequence according to SEQ ID NO: 1

to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30, and the equivalent positions within SEQ ID NO: 5 to SEQ ID NO: 28. Said oligonucleotides may also be present in the form of peptide nucleic acids. The non-hybridized amplicates are then removed. The hybridized amplicates are then detected. In this context, it is preferred that labels attached to the amplicates are identifiable at each position of the solid phase at which an oligonucleotide sequence is located.

In yet a further embodiment of the method, the genomic methylation status of the CpG positions may be ascertained by means of oligonucleotide probes that are hybridised to the bisulfite treated DNA concurrently with the PCR amplification primers (wherein said primers may either be methylation specific or standard).

A particularly preferred embodiment of this method is the use of fluorescence-based Real Time Quantitative PCR (Heid et al., *Genome Res.* 6:986-994, 1996; also see United States Patent No. 6,331,393) employing a dual-labeled fluorescent oligonucleotide probe (TaqMan™ PCR, using an ABI Prism 7700 Sequence Detection System, Perkin Elmer Applied Biosystems, Foster City, California). The TaqMan™ PCR reaction employs the use of a nonextendible interrogating oligonucleotide, called a TaqMan™ probe, which, in preferred embodiments, is designed to hybridize to a GpC-rich sequence located between the forward and reverse amplification primers. The TaqMan™ probe further comprises a fluorescent "reporter moiety" and a "quencher moiety" covalently bound to linker moieties (e.g., phosphoramidites) attached to the nucleotides of the TaqMan™ oligonucleotide. For analysis of methylation within nucleic acids subsequent to bisulfite treatment, it is required that the probe be methylation specific, as described in United States Patent No. 6,331,393, (hereby incorporated by reference in its entirety) also known as the MethyLight™ assay. Variations on the TaqMan™ detection methodology that are also suitable for use with the described invention include the use of dual-probe technology (Lightcycler™) or fluorescent amplification primers (Sunrise™ technology). Both these techniques may be adapted in a manner suitable for use with bisulfite treated DNA, and moreover for methylation analysis within CpG dinucleotides.

A further suitable method for the use of probe oligonucleotides for the assessment of methylation by analysis of bisulfite treated nucleic acids. In a further preferred embodiment of the method, the *fifth step* of the method comprises the use of template-directed oligonucleotide extension, such as MS-SNuPE as described by Gonzalzo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997.

In yet a further embodiment of the method, the *fifth step* of the method comprises sequencing and subsequent sequence analysis of the amplificate generated in the *third step* of the method (Sanger F., et al., *Proc Natl Acad Sci USA* 74:5463-5467, 1977).

Best mode

In the most preferred embodiment of the method the nucleic acids according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 are isolated and treated according to the first three steps of the method outlined above, namely:

- a. obtaining, from a subject, a biological sample having subject genomic DNA;
- b. extracting or otherwise isolating the genomic DNA;
- c. treating the genomic DNA of b), or a fragment thereof, with one or more reagents to convert cytosine bases that are unmethylated in the 5-position thereof to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridization properties;

and wherein the subsequent amplification of d) is carried out in a methylation specific manner, namely by use of methylation specific primers or *blocking oligonucleotides*, and further wherein the detection of the amplicates is carried out by means of a real-time detection probes, as described above.

Wherein the subsequent amplification of d) is carried out by means of methylation specific primers, as described above, said methylation specific primers comprise a sequence having a length of at least 9 nucleotides which hybridizes to a pretreated nucleic acid sequence according to one of SEQ ID NO: 5 to SEQ ID NO: 28 and sequences complementary thereto, wherein the base sequence of said oligomers comprises at least one CpG dinucleotide. Step e) of the method, namely the detection of

the specific amplicates indicative of the methylation status of one or more CpG positions according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 is carried out by means of real-time detection methods as described above.

In an alternative most preferred embodiment of the method the subsequent amplification of d) is carried out in the presence of *blocking oligonucleotides*, as described above. Said *blocking oligonucleotides* comprising a sequence having a length of at least 9 nucleotides which hybridizes to a pretreated nucleic acid sequence according to one of SEQ ID NO: 5 to SEQ ID NO: 28 and sequences complementary thereto, wherein the base sequence of said oligomers comprises at least one CpG, TpG or CpA dinucleotide. Step e) of the method, namely the detection of the specific amplicates indicative of the methylation status of one or more CpG positions according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 is carried out by means of real-time detection methods as described above.

14 14 14

Diagnostic and/or Prognostic Assays for prostate cell proliferative disorders

The present invention enables diagnosis of events which are disadvantageous to patients or individuals in which important genetic and/or epigenetic parameters within one or more of the genes GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions including genomic sequences according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 may be used as markers. Said parameters obtained by means of the present invention may be compared to another set of genetic and/or epigenetic parameters, the differences serving as the basis for a diagnosis and/or prognosis of events which are disadvantageous to patients or individuals.

Specifically, the present invention provides for diagnostic and/or prognostic cancer assays based on measurement of differential methylation of one or more CpG dinucleotide sequences of the genes GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions including genomic sequences according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID

NOs: 29 & 30, or of subregions thereof that comprise such a CpG dinucleotide sequence. Typically, such assays involve obtaining a tissue sample from a test tissue, performing an assay to measure the methylation status of at least one of one or more CpG dinucleotide sequences of the genes GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions including genomic sequences according to SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30 derived from the tissue sample, relative to a control sample, or a known standard and making a diagnosis or prognosis based thereon.

In particular preferred embodiments, inventive oligomers are used to assess the CpG dinucleotide methylation status, such as those based on SEQ ID NO: 1 to SEQ ID NO: 30, or arrays thereof, as well as in kits based thereon and useful for the diagnosis and/or prognosis of prostate cell proliferative disorders.

Kits

Moreover, an additional aspect of the present invention is a kit comprising, for example: a bisulfite-containing reagent; a set of primer oligonucleotides containing at least two oligonucleotides whose sequences in each case correspond, are complementary, or hybridize under stringent or highly stringent conditions to a 16-base long segment of one or more of the genes GSTP1, HISTONE H4, PROSTAGLANDIN E2 RECEPTOR, LIM DOMAIN KINASE 1, SEQ ID NO: 29 & ORPHAN NUCLEAR RECEPTOR NR5A2 and their regulatory regions including genomic and/or treated sequences according to SEQ ID NO: 1 to SEQ ID NO: 30; oligonucleotides and/or PNA-oligomers; as well as instructions for carrying out and evaluating the described method. In a further preferred embodiment, said kit may further comprise standard reagents for performing a CpG position-specific methylation analysis, wherein said analysis comprises one or more of the following techniques: MS-SNuPE, MSP, MethyLight™, HeavyMethyl™, COBRA, and nucleic acid sequencing. However, a kit along the lines of the present invention can also contain only part of the aforementioned components.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following example serves only to illustrate the invention and is not intended to limit the invention within the principles and scope of the broadest interpretations and equivalent configurations thereof.

EXAMPLES

The objective of the following study was to analyze the methylation status of prostate cancer markers in different body fluid samples in order to identify the preferred choice of body fluid (urine or serum) for testing and the preferred marker, markers or combinations of markers. The study was run on matched serum and urine sediment samples from 80 patients with an average age of 65 and representative of a number of racial types (caucasian, african american etc..). In each case, genomic DNA was analyzed using the HeavyMethyl or MSP technique after bisulfite conversion.

Urine Sediment was prepared for analysis and bisulphite treated according to the following:

- 200 ul sediment samples were purified using the Magnapure DNA Isolation Kit 1 with a 100 ul elution volume.
- 5 ul HD6 PCR was carried out on the Magnapure Eluate , in order to determine DNA concentration
- 100 ul of the DNA solution was treated using a proprietary bisulfite treatment technique
- 10 ul C3 bisulfite specific quantitative PCR
- 5 ul Merck sulfite test

Serum was prepared for analysis and bisulphite treated according to the following:

- 1 mL serum samples were purified using the Magnapure DNA Large Volume Total nucleic acid with a 100 ul elution volume.
- 5 ul HD6 PCR on Magnapure Eluate - To determine DNA concentration

- 100 ul of the DNA solution was treated using a proprietary bisulfite treatment technique
- 10 ul C3 bisulfite specific quantitative PCR
- 5 ul Merck sulfite test

Single PCR runs were performed on 10 ul of bisulfite treated DNA per sample for each of the markers as described below.

Heavy Methyl Assay of the GSTP1 gene

In the following analysis the methylation status of the gene GSTP1 was analysed by means of methylation specific amplification using the primers according to Table 1 (below).

The sequence of interest is amplified by means of methylation specific primers and a blocker oligonucleotide in order to minimise the unspecific amplification of non methylated DNA. The amplificate is then detected by means of methylation specific Lightcycler probes.

Table 1: Oligonucleotides for MSP - Lightcycler analysis of GSTP1.

SEQ ID NO:	Sequence	Type
31	gggattatTTTTATAaggTt	primer
32	ctctaaaccccatcccc	primer
33	cccatccccaaaaacacaaaccac	blocker
34	CGtCGtCGtAGTtTTCGtt-fluo	probe
35	red640-tAGTGAGTACGCGCGGtt-pho	probe

Reaction conditions:

PCR program

denat at 95°C

95°C

10min

50 cycles:

ramp

denat at 95°C 10 sec (1°C/s)

annealing 56°C 30 sec (1°C/s) detection
 extension 72°C 10 sec (1°C/s)

MSP analysis of the gene HISTONE H4.

In the following analysis the methylation status of the gene HISTONE H4 was analysed by means of methylation specific amplification using the primers according to Table 2 (below).

The sequence of interest is amplified by means of methylation specific primers, the amplificate is then detected by means of methylation specific Taqman probes.

Table 2: Oligonucleotides for MSP - Taqman analysis of HISTONE H4.

SEQ ID NO :	Sequence	Type
36	accgaaaatacgcttcacg	primer
37	gcgttatcgtaaagtattgcgc	primer
38	/56-FAM/cgcgacgaacaaaacgccg/3BHQ_1/	probe

Reaction Conditions:

PCR program

denat at 95°C

95°C 10min

50 cycles: ramp

denat at 95°C 10 sec (20°C/s)

annealing 60°C 45 sec (20°C/s) detection

MSP analysis of the gene PROSTAGLANDIN E2 RECEPTOR.

In the following analysis the methylation status of the gene PROSTAGLANDIN E2 RECEPTOR was analysed by means of methylation

specific amplification using the primers according to Table 3 (below) .:

The sequence of interest is amplified by means of methylation specific primers, the amplificate is then detected by means of methylation specific Taqman probes.

Table 3: Oligonucleotides for MSP - Taqman analysis of PROSTAGLANDIN E2 RECEPTOR

SEQ ID NO :	Sequence	Type
39	cgcgctactccgcataca	primer
40	gaggtaatcgaggcggtcg	primer
41	/56-FAM/cgccaattcatacgccgcacc/3BHQ_1/	probe

PCR program

denat at 95°C

95°C 10min

50 cycles: ramp

denat at 95°C 10 sec (20°C/s)

annealing 60°C 45 sec (20°C/s) detection

MSP analysis of the gene ORPHAN NUCLEAR RECEPTOR NR5A2.

In the following analysis the methylation status of the gene ORPHAN NUCLEAR RECEPTOR NR5A2 was analysed by means of methylation specific amplification using the primers according to Table 3 (below) .

The sequence of interest is amplified by means of methylation specific primers, the amplificate is then detected by means of methylation specific Taqman probes.

Table 4: Oligonucleotides for MSP - Taqman analysis of ORPHAN NUCLEAR RECEPTOR NR5A2.

SEQ ID NO :	Sequence	Type
42	ttgtggttcgggaagagac	primer
43	tcccgaactcttcgatcg	primer
44	aactacgcgcaaaccgcga	probe

PCR program

denat at 95°C

95°C

10min

50 cycles:

ramp

denat at 95°C 10 sec (20°C/s)

annealing 60°C 45 sec (20°C/s) detection

Marker Analysis

Results were analyzed qualitatively by scoring amplification as +/- and quantitatively by determining the percentage of methylated DNA as a fraction of total DNA calculated using the C3 bisulfite specific PCR. To measure total methylated DNA, a 100% methylated standard (chemicon SSS1 treated DNA) standard curve was included in each assay.

Results

For each marker a Receiver Operating Characteristic curve (ROC curve) of the assay was determined. A ROC is a plot of the true positive rate against the false positive rate for the different possible cutpoints of a diagnostic test. It shows the tradeoff between sensitivity and specificity depending on the selected cutpoint (any increase in sensitivity will be accompanied by a decrease in specificity). The area under an ROC curve (AUC) is a measure for the accuracy of a diagnostic test (the larger the area the better, optimum is 1, a random test would have a ROC

curve lying on the diagonal with an area of 0.5; for reference: J.P. Egan. Signal Detection Theory and ROC Analysis, Academic Press, New York, 1975).

AUC results:

Serum:

Marker: HeavyMethyl GSTP1

AUC: 0.51

Marker: MSP HISTONE H4

AUC: 0.59

Marker: MSP PROSTAGLANDIN E2 RECEPTOR

AUC: 0.52

Marker: MSP ORPHAN NUCLEAR RECEPTOR NR5A2

AUC: 0.50

Urine:

Marker: HeavyMethyl GSTP1

AUC: 0.58

Marker: MSP HISTONE H4

AUC: 0.5

Marker: MSP PROSTAGLANDIN E2 RECEPTOR

AUC: 0.49

Marker: MSP ORPHAN NUCLEAR RECEPTOR NR5A2

AUC: 0.56

In order to provide an accurate detection of prostate cancer it is preferred that a combined analysis of multiple markers is carried out (i.e. a gene panel). For analysis of urine based samples the most preferred combination of markers is GSTP1 , PROSTAGLANDIN E2 RECEPTOR & ORPHAN NUCLEAR RECEPTOR NR5A2 with a sensitivity of 0.37 and a specificity of 0.72.

For analysis of serum based samples the most preferred combination of markers is GSTP1 , HISTONE H4 & ORPHAN NUCLEAR RECEPTOR NR5A2 with a sensitivity of 0.35 and a specificity of 0.75.

MSP analysis of the genes according to Table 1.

In the following analysis the methylation status of a selection of the genes according to Table 5 were analysed by means of MSP TaqMan assays using the primers and probes according to Table 6 (below).

The study was run on 10 samples from prostate carcinoma tissue, 10 from benign prostate hyperplasia tissue and 5 normal prostate tissue samples. Genomic DNA was analyzed using the MSP technique after bisulfite conversion. Total genomic DNA of all samples was bisulfite treated converting unmethylated cytosines to uracil. Methylated cytosines remained conserved. Bisulfite treatment was performed with minor modifications according to the protocol described in Olek et al. (1996).

The sequence of interest was then amplified by means of primers specific for bisulfite treated DNA and the amplicates were detected by means of TaqMan probes using TaqMan and/or Lightcycler platforms. Results are shown in table 7 below.

By combining SEQ ID NO: 29 with SEQ ID NO: 3 a sensitivity of 58% and a specificity of 92% was achieved.

Reagents and cycling conditions were as follows:

SEQ ID NO: 3

Reagent	Final Conc.
Reverse Primer	10 ⁻⁴ M
Reverse Primer	10 ⁻⁴ M
Probe	10 ⁻⁴ M
MgCl ₂	2.5 mM
Tris-HCl	10 mM
Water	10 mM
Template	10 ⁻⁴ M
Total	10 ⁻⁴ M

Temperature (C)	Time (sec)	Rate (C/sec)	Collector F	# of Cycles
Dehydration				1
95	30	120		
Annealing				6
95	10			
65	45			
Cool				1
40	300	20		

SEQ ID NO: 2

Reagent	Vol Volume (μ l)
Forward Primer	35.0
Reverse Primer	35.1
Probe	2.8
Water	12.2
Template	245.0
Total	500.0

Temperature (C)	Time (sec)	# of Cycles
Dehydration		
95	30	
Annealing		50
95		
60	25	

SEQ ID NO: 30

Reagent	Vol Volume (μ l)
Forward Primer	17.5
Reverse Primer	17.5
Probe	0.4
MgCl ₂	1.5
Tris-Cl	1.0
Water	1.0
Template	1.0

Seq. ID NO.	29
-------------	----

Sample Name	Time (sec)	Ratio (C/50)	Collect	Cycles
Blank	10	1.0		1
10	10	1.0		1
20	10	1.0		1
30	10	1.0		1
40	10	1.0		1
50	10	1.0		1
60	10	1.0		1
70	10	1.0		1
80	10	1.0		1
90	10	1.0		1
100	10	1.0		1

SEQ ID NO: 29

Reagent	Volume (µl)
Forward Primer	1.0
Reverse Primer	1.0
Probe	1.0
Water	1.0
Template	1.0
Total	5.0

Sample Name	Time (sec)	Cycles
Blank	10	1
10	10	1
20	10	1
30	10	1
40	10	1
50	10	1
60	10	1
70	10	1
80	10	1
90	10	1
100	10	1

Table 5: Genes & Sequences according to the invention

Gene Name	Genbank Ref. Seq.	Genomic sequence SEQ ID NO:	Treated sequences (methylated) SEQ ID NO:	Treated sequences (unmethylated) SEQ ID NO:
GSTP1	NM_000852	1	5 & 6	13 & 14
HISTONE H4	NM_003495	2	7 & 8	15 & 16

PROSTAGLANDIN E2 RECEPTOR	NM_000958	3	9 & 10	17 & 18
ORPHAN NUCLEAR RECEPTOR NR5A2	NM_003822	4	11 & 12	19 & 20
2308bp non-coding region of 19p13.2*	Contig AC020947.6.1.36838	29	21 & 22	25 & 26
LIM DOMAIN KINASE 1	NM_002314	30	23 & 24	27 & 28

*the sequence is located within the promoter regions of
ENSESTG00002947491 & upstream of upstream of ACP5.

Table 6

Genomic sequence SEQ ID NO:	Forward primer SEQ ID NO:	Reverse primer SEQ ID NO:	Probe SEQ ID NO:
3	45	46	47
2 (assay 4)	48	49	50
2 (assay 1)	51	52	53
30	54	55	56
29	57	58	59

Table 7

Genomic sequence SEQ ID NO:	Sensitivity (%)	Specificity
3	70	88
2 (assay 1)	60	94
30	80	80
29	80	79
2 (assay 4)	60	98

10-05-2004

We claim:

1. A method for detecting, or for detecting and distinguishing between or among prostate cell proliferative disorders in a subject with a sensitivity of greater than 30% and a specificity of greater than 65%, said method comprising analysing the methylation pattern of a target nucleic acid comprising one or a combination of sequences taken from the group consisting of SEQ ID Nos: 1-4, 29 & 30 by contacting at least one of said target nucleic acids in a biological sample obtained from said subject with at least one reagent, or series of reagents that distinguishes between methylated and non-methylated CpG dinucleotides.
2. The method of claim 1, wherein prostate carcinoma is distinguished from at least one condition selected from the group consisting of prostate adenoma, normal prostate tissue, non-prostate tissues and non-prostate cell proliferative disorders.
3. A method according to claim 1, comprising:
 - obtaining, from a subject, a biological sample having subject genomic DNA;
 - contacting the genomic DNA, or a fragment thereof, with one reagent or a plurality of reagents for distinguishing between methylated and non methylated CpG dinucleotide sequences within at least one target sequence of the genomic DNA, or fragment thereof, wherein the target sequence comprises, or hybridizes under stringent conditions to, at least 16 contiguous nucleotides of a sequence taken from the group consisting of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOs: 29 & 30, said contiguous nucleotides comprising at least one CpG dinucleotide sequence; and
 - determining, based at least in part on said distinguishing, the methylation state of at least one target CpG dinucleotide sequence, or an average, or a value reflecting an average methylation state of a plurality of target CpG dinucleotide sequences, whereby

detecting, or detecting and distinguishing between or among prostate cell proliferative disorders with a sensitivity of greater than 30% and a specificity of greater than 65% is, at least in part, afforded.

4. The method of claim 3, wherein distinguishing between methylated and non methylated CpG dinucleotide sequences within the target sequence comprises converting unmethylated cytosine bases within the target sequence to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridization properties.
5. The method of claim 3, wherein distinguishing between methylated and non methylated CpG dinucleotide sequences within the target sequence(s) comprises methylation state-dependent conversion or non-conversion of at least one CpG dinucleotide sequence to the corresponding converted or non-converted dinucleotide sequence within a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and contiguous regions thereof corresponding to the target sequence.
6. The method of claim 3, wherein the biological sample is selected from the group consisting of cell lines, histological slides, biopsies, paraffin-embedded tissue, bodily fluids, ejaculate, urine, blood, and combinations thereof.
7. The method of claim 3, wherein distinguishing between methylated and non methylated CpG dinucleotide sequences within the target sequence comprises use of at least one nucleic acid molecule or peptide nucleic acid (PNA) molecule comprising, in each case a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof.

8. The method of claim 7, wherein the contiguous sequence comprises at least one CpG, TpG or CpA dinucleotide sequence.
9. The method of claim 7, comprising use of at least two such nucleic acid molecules, or peptide nucleic acid (PNA) molecules.
10. The method of claim 7, comprising use of at least two such nucleic acid molecules, or peptide nucleic acid (PNA) molecules as primer oligonucleotides for the amplification of a sequences selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, sequences complementary thereto, and regions thereof that comprise, or hybridize under stringent conditions to the primers.
11. The method of claim 9, comprising use of at least four such nucleic acid molecules, or peptide nucleic acid (PNA) molecules.
12. A method for detecting, or detecting and distinguishing between or among prostate cell proliferative disorders in a subject, comprising:
 - a. obtaining, from a subject, a biological sample having subject genomic DNA;
 - b. extracting or otherwise isolating the genomic DNA;
 - c. treating the genomic DNA of b), or a fragment thereof, with one or more reagents to convert cytosine bases that are unmethylated in the 5-position thereof to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridization properties;
 - d. contacting the treated genomic DNA, or the treated fragment thereof, with an amplification enzyme and at least two primers comprising, in each case a contiguous sequence of at least 9 nucleotides that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, wherein the

treated genomic DNA or the fragment thereof is either amplified to produce at least one amplificate, or is not amplified; and

e) determining, based on a presence or absence of, or on a property of said amplificate, the methylation state of at least one CpG dinucleotide of a sequence selected from the group consisting SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 , or an average, or a value reflecting an average methylation state of a plurality of CpG dinucleotides of a sequence selected from the groups consisting of SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 , whereby at least one of detecting, or detecting and distinguishing between prostate cell proliferative disorders with a sensitivity of greater than 30% and a specificity of greater than 65% is at least in part, afforded.

13. The method of claim 12, wherein treating the genomic DNA, or the fragment thereof in c), comprises use of a reagent selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof.
14. The method of claim 12, wherein contacting or amplifying in d) comprises use of at least one method selected from the group consisting of: use of a heat-resistant DNA polymerase as the amplification enzyme; use of a polymerase lacking 5'-3' exonuclease activity; use of a polymerase chain reaction (PCR); generation of a amplificate nucleic acid molecule carrying a detectable labels; and combinations thereof.
15. The method of claim 14, wherein the detectable amplificate label is selected from the label group consisting of: fluorescent labels; radionuclides or radiolabels; amplificate mass labels detectable in a mass spectrometer; detachable amplificate fragment mass labels detectable in a mass spectrometer; amplificate, and detachable amplificate fragment mass labels having a single-positive or single-negative net

charge detectable in a mass spectrometer; and combinations thereof.

16. The method of claim 12, wherein the biological sample obtained from the subject is selected from the group consisting of cell lines, histological slides, biopsies, paraffin-embedded tissue, bodily fluids, ejaculate, urine, blood, and combinations thereof.
17. The method of claim 12, wherein prostate carcinoma is distinguished from at least one condition selected from the group consisting of prostate adenoma, inflammatory prostate tissue, prostate adenomas with grade 2 dysplasia less than 1 cm, prostate adenomas with grade 3 dysplasia equal to or greater than 1 cm in size, normal prostate tissues, non-prostate normal tissue, body fluids, and non-prostate cancer tissue.
18. The method of claim 12, further comprising in step d) the use of at least one nucleic acid molecule or peptide nucleic acid molecule comprising in each case a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, wherein said nucleic acid molecule or peptide nucleic acid molecule suppresses amplification of the nucleic acid to which it is hybridized.
19. The method of claim 18, wherein said nucleic acid molecule or peptide nucleic acid molecule is in each case modified at the 5'-end thereof to preclude degradation by an enzyme having 5'-3' exonuclease activity.
20. The method of claim 18, wherein said nucleic acid molecule or peptide nucleic acid molecule is in each case lacking a 3' hydroxyl group.

21. The method of claim 18, wherein the amplification enzyme is a polymerase lacking 5'-3' exonuclease activity.
22. The method of claim 12, wherein determining in e) comprises hybridization of at least one nucleic acid molecule or peptide nucleic acid molecule in each case comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof.
23. The method of claim 22, wherein at least one such hybridizing nucleic acid molecule or peptide nucleic acid molecule is bound to a solid phase.
24. The method of claim 22, wherein a plurality of such hybridizing nucleic acid molecules or peptide nucleic acid molecules are bound to a solid phase in the form of a nucleic acid or peptide nucleic acid array selected from the array group consisting of linear or substantially so, hexagonal or substantially so, rectangular or substantially so, and combinations thereof.
25. The method of claim 22, further comprising extending at least one such hybridized nucleic acid molecule by at least one nucleotide base.
26. The method of claim 12, wherein determining in e), comprises sequencing of the amplificate.
27. The method of claim 12, wherein contacting or amplifying in d), comprises use of methylation-specific primers.
28. The method of claim 12 comprising in d) using primer oligonucleotides comprising one or more CpG; TpG or CpA dinucleotides; and further comprising in e) the use of at least one method selected from the group

consisting of: hybridizing in at least one nucleic acid molecule or peptide nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof; hybridizing at least one nucleic acid molecule that is bound to a solid phase and comprises a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof; hybridizing at least one nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, and extending at least one such hybridized nucleic acid molecule by at least one nucleotide base; and sequencing in e) of the amplificate.

29. The method of claim 12 comprising in d) use of at least one nucleic acid molecule or peptide nucleic acid molecule comprising in each case a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, wherein said nucleic acid molecule or peptide nucleic acid molecule suppresses amplification of the nucleic acid to which it is hybridized; and further comprising in e) the use of at least one method selected from the group consisting of: hybridizing in at least one nucleic acid molecule or peptide nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof; hybridizing at least

30. The method of claim 12, comprising in d) amplification by primer oligonucleotides comprising one or more CpG; TpG or CpA dinucleotides and further comprising in e) hybridizing at least one detectably labeled nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28.
31. The method of claim 12, comprising in d) the use of at least one nucleic acid molecule or peptide nucleic acid molecule comprising in each case a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and complements thereof, wherein said nucleic acid molecule or peptide nucleic acid molecule suppresses amplification of the nucleic acid to which it is hybridized, and further comprising in e) hybridizing at least one detectably labeled nucleic acid molecule comprising a contiguous sequence at least 9 nucleotides in length that is complementary to, or hybridizes under

moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28.

32. A treated nucleic acid derived from genomic SEQ ID NO: 1 to SEQ ID NO: 4 and SEQ ID NOS: 29 & 30 , wherein the treatment is suitable to convert at least one unmethylated cytosine base of the genomic DNA sequence to uracil or another base that is detectably dissimilar to cytosine in terms of hybridization.
33. A nucleic acid, comprising at least 16 contiguous nucleotides of a treated genomic DNA sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28, and sequences complementary thereto, wherein the treatment is suitable to convert at least one unmethylated cytosine base of the genomic DNA sequence to uracil or another base that is detectably dissimilar to cytosine in terms of hybridization.
34. The nucleic acid of claims 32 and 33 wherein the contiguous base sequence comprises at least one CpG, TpG or CpA dinucleotide sequence.
35. The nucleic acid of claims 32 and 33 wherein the treatment comprises use of a reagent selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof .
36. An oligomer, comprising a sequence of at least 9 contiguous nucleotides that is complementary to, or hybridizes under moderately stringent or stringent conditions to a treated genomic DNA sequence selected from the group consisting of SEQ ID NO: 5 to SEQ ID NO: 28.
37. The oligomer of Claim 36, comprising at least one CpG , CpA or TpG dinucleotide sequence.

38. A set of oligomers, comprising at least two oligonucleotides according, in each case, to any one of Claims 37.
39. Use of a set of oligomers according, in each case, to any one of Claims 36 through 38, as probes for determining at least one of a cytosine methylation state, or a single nucleotide polymorphism (SNP) of a sequence selected from the group consisting of SEQ ID NO: 1 to 4, 29 & 30 and sequences complementary thereto.
40. A kit useful for detecting, or for detecting distinguishing between or among prostate cell proliferative disorders of a subject, comprising:
- at least one of a bisulfite reagent, or a methylation-sensitive restriction enzyme; and
 - at least one nucleic acid molecule or peptide nucleic acid molecule comprising, in each case a contiguous sequence at least 9 nucleotides that is complementary to, or hybridizes under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID 5 to SEQ ID NO 28, and complements thereof
41. The kit of claim 40, further comprising standard reagents for performing a methylation assay selected from the group consisting of MS-SNuPE, MSP, MethyLight, HeavyMethyl, COBRA, nucleic acid sequencing, and combinations thereof.
42. The method of any one of claims 1, 12 or 3 comprising use of the kit according to claim 41.
43. Use of a nucleic acid according to claims 32 through 35, an oligomer according to any one of claims 36 through 37, a set of oligonucleotides according to claim 38 and a kit according to claims 41 and 42 for the detection of,

detection and differentiation between or among subclasses
of prostate cell proliferative disorders.

10-05-2004

Abstract

The invention provides methods, nucleic acids and kits for detecting, or for detecting and distinguishing between or among prostate cell proliferative disorders. The invention discloses genomic sequences the methylation patterns of which have utility for the improved detection of and differentiation between said class of disorders, thereby enabling the improved diagnosis and treatment of patients.

<110> Epigenomics AG

<120> METHODS AND NUCLEIC ACIDS FOR THE ANALYSIS OF CpG DINUCLEOTIDE METHYLATION STATUS ASSOCIATED WITH THE DEVELOPMENT OF PROSTATE CANCER.

<160> 34

EPO-BERLIN

<210> 1

<211> 2501

<212> DNA

<213> Homo Sapiens

10 -05- 2004

<400> 1

ttgttgtaga	gaatattttca	tcacccaggt	attatgccga	gtacccaata	gttctctttt	60
ctgctcctct	ccttcctccc	atcctgcacc	ctggagtcac	ccacagtgtc	tggtgtttcc	120
ttgttttgtg	tataagttct	catcatttag	ctcccactta	caagtggaga	catccagtat	180
ttggattttct	gttcctgcat	tagtttgcta	aggataatag	cctctagctc	catccatgtt	240
cccacaaaag	acatgatcta	gttcttttta	atggctgcat	taaatgaagt	tttaaagata	300
caacataaac	accaacctct	tccccaccac	aaaaatccct	tgctgaattt	gattacactt	360
aaattaacga	gttttgtttc	atgaaagact	ccttggacaa	acttgacagt	tgatgggaata	420
ggagaagctg	tctgtcatgt	ctaaagccaa	caagagatca	atatctagaa	taaatggaga	480
tctgcaaatc	aacagaaagt	aggcagcaaa	gccaaagaaa	atagcctaag	gcacagccac	540
taaaaggaac	gtgatcatgt	cctttgcagg	gacatgggtg	gagctggaag	ccgttagcct	600
cagcaaaactc	acacaggaac	agaaaaccag	cgagaccgca	tggtctcact	tataagtggg	660
agctgaacaa	tgagaacaca	tggtcacatg	gcggcgatca	acacacactg	gtgcctgttg	720
agcggggtgc	tggggaggga	gagtaccagg	aagaatagct	aagggtact	gggcttaata	780
cctgggtgat	gggatgatct	gtacagcaaa	ccatcatggc	gcacacacct	atgtaacaaa	840
cctgcacatc	ctctacatgt	accccagaac	ttcaaatata	agttggacgg	ccaggcgtgg	900
tggtcacgcg	ctgtaatccc	agcacttttg	gaagccgagg	cgtgcagatc	acctaaggct	960
aggagttcga	gaccagcccg	gccaacatgg	tgaacccccg	tctctactaa	aaatacaaaa	1020
atcagccaga	tgtggcacgc	acctataatt	ccacctaact	gggaggctga	agcagaattg	1080
cttgaacccg	agaggcggag	gttgacagtga	gccgccgaga	tgcgccact	gcactccagc	1140
ctggggccaca	gcgtgagact	acgtcataaa	ataaaataaa	ataacacaaa	ataaaataaa	1200
ataaaataaaa	ataaaataaa	ataaaataaa	ataaaataaa	ataaaaaaat	aaaaataaat	1260
aaaataaaaat	aaagcaattt	cctttcctct	aagcggcctc	cacccctctc	ccctgccctg	1320
tgaagcgggt	gtgcaagctc	cgggatcgca	gcggtcttag	ggaatttccc	cccgcgatgt	1380
cccggcgcgc	cagttcgctg	cgcacacttc	gctgcggctc	tcttctctgt	gtctgtttac	1440
tccctaggcc	ccgctgggga	cctgggaaag	agggaaaggc	ttccccggcc	agctgcgcgg	1500
cgactccggg	gactccaggg	cgccctctct	cgcccgacgc	ccggggtgca	gcggccgcgc	1560
gggctggggc	ggcggggagt	ccgcgggacc	ctccagaaga	gcggccggcg	ccgtgactca	1620
gcactggggc	ggagcggggc	gggaccaccc	ttataaggct	cgagggccgc	gaggccttcg	1680
ctggagtttc	gccgccgcag	tcttcgccac	cagttagtac	gcgcggcccc	cgtccccggg	1740
gatggggctc	agagctccca	gcatggggcc	aaccgcagc	atcaggcccc	ggctcccggc	1800
agggctcctc	gccacctcgc	agaccgggga	cgggggccta	ggggacccag	gacgtcccca	1860
gtgccgttag	cggttttcag	ggggcccgga	gcgcctcggg	gagggatggg	accccggggg	1920
cgggggagggg	gggcagactg	cgctcacgcg	gccttggcat	cctcccccg	gctccagcaa	1980
actttttctt	gttcgctgca	gtgccgccct	acaccgtggg	ctatttccca	gttcgaggta	2040
ggagcatgtg	tctggcaggg	aagggaggga	ggggctgggg	ctgcagccca	cagcccctcg	2100
cccaccggga	gagatccgaa	cccccttata	cctccgtcgt	gtggctttta	ccccgggcct	2160
ccttctctgt	ccccgcctct	cccgccatgc	ctgctccccg	ccccagtgtt	gtgtgaaatc	2220
ttcggaggaa	cctgtttccc	tggtccctcc	ctgcactcct	gacccctccc	cggtgtgtcg	2280
cgaggcggag	tccggccggg	ccccacatct	cgtacttctc	cctccccgca	ggccgctgcg	2340
cggcctctcg	catgctgctg	gcagatcagg	cccagagctg	gaaggaggag	gtggtgaccg	2400
tgagagcgtg	gcaggagggc	tactcaaaag	cctcctgcgt	aagtgaccat	gcccggggcaa	2460
ggggaggggg	tgctgggcct	tagggggctg	tgactaggat	c		2501

<210> 2

<211> 2501

<212> DNA

<213> Homo Sapiens

<400> 2

tttgcaaatg	gagacatctt	cattattcct	atagtatcat	atgtttttta	agtttgtact	60
cacacttttg	gtgataaatg	aaggacaaga	tccttcccta	tccttgtgag	gatgactaca	120
gcatgactgg	atgggcttgc	tatgattttt	atctttccct	gtgttctcac	taccgtttta	180
ttaatctcag	ttctttttca	cagggtagca	cagaatttaa	ctagcagaaa	gagatccagc	240
catgtagacc	agagatttgt	ctaagtgcag	gcatgtaaga	atcaggaagg	aaagtttttt	300
gttttaaatac	caacagggttc	cttccttaaa	gcaattatta	tttttcaaat	ctaaccacaca	360

aggtgatagt	atccttaaac	caattaaatc	agaatctcgg	gttggataac	ctcaaatacg	420
acttattagc	acttcccatt	aatcactggg	ccttcaggcc	tttaagttta	cttactagga	480
atctcacttt	taataccatc	ttatcaactt	cagttgtaaa	taagagaaca	ctcaaaggct	540
gaggaattct	cagcggtaaa	gctctgcccc	cgtaagttaa	caaaggataa	gttagtcttt	600
gttgatgaca	ctttgttgta	ctgataagct	acgtatttct	actcaaggat	tcaaattctc	660
acctttctca	agaattgggc	caaaaccgat	aaactaaact	tatttacggg	ccactgatta	720
aaggttggtg	cataataagt	tcttgctatg	ttcagcagtt	ggattcacag	cgccagaaac	780
ctataactgc	ttgactttcc	tccccactac	actgcgaaaa	ttgcccctta	aatgtaacta	840
accctaaaac	ctcaacagta	tcgtggccag	gcgtggtggc	tcactactgt	aataccaaca	900
ttaggcatag	gcgaggggat	tgaggccagg	atatcgaaac	tagcctggga	aacacacgga	960
gacccggtct	ttggaaaaat	aattagcctt	gcgtggtggt	gggcgcgagg	ttccggctaa	1020
tcgggaggct	acagtgaagg	atgatgacac	tgactacag	tctgcgcgac	ggcccatgtc	1080
agtaagctct	ggagcacctg	aaacaagttg	tgttgggtat	tttatttact	ggagagcgat	1140
tagtgactga	tgcctactta	cagcgactag	agacgcagtc	tccgatagca	gcacaaactc	1200
agcaggcgcg	aacaaatggt	aaagagaaac	tgggcaaaac	agcatcacgg	ctcctcagct	1260
gagaaagtgg	gggccctaaa	aagggccttt	tgttgataga	aagggacgct	caaccaccga	1320
aaccgtagag	ggtgcggccc	tggcgcttga	gcgcgtagac	cacatccatg	gcggtgaccg	1380
tcttgcgctt	ggcgtgctct	gtataggtca	cggcgctccc	gatcacgttc	tccaggaaca	1440
ccttcagcac	cccgcgagtc	tcttcgtaga	tgaggccgga	gatgcgcttc	acgccgcgcg	1500
ggcgagcaag	gcgcccgatg	gccggcttgg	tgatgccctg	gatattgtcg	cgcagtactt	1560
tacggtggcg	ccttagcgcg	cctttgccaa	gaccttccc	gcctttgccg	cgccagaca	1620
tgacgagcaa	gaggagtctc	acccaacgct	ttgtgaggac	tctggcctga	ggcagcgctt	1680
ttatacgaca	gttgggcgac	cgaactgaga	acctgaaaga	agtcggcggg	aagtcccgcg	1740
ccggtggggg	aggggaaatc	taaagggcca	aaccgaaata	gggggaaaaa	aaaagcgagc	1800
ttcttgtttc	cgtgtttctg	atthttgtaac	gtgcatagta	ttttgttacc	acgttatgag	1860
gctttaaaaa	attgcttttg	aacgcagaag	atatacatca	atactgtggg	aaatacaaga	1920
aaggacaaga	aattaagaaa	ctacaatggt	atcccatcac	acaggctagt	taatcatgta	1980
ttttgcagag	cagtgcaca	tatttttcca	agaaaatgta	tacagtgttg	tatatggagt	2040
tttgtaacct	ccttatattg	attataatth	aaccaatttc	tattaaagag	ataaaaagtga	2100
tgttttggtg	tctatgtttc	ttaggaatta	tcaatagtta	taatcagttc	cccagcaatt	2160
ttttaatcgg	ctgtatttta	tttccacatt	gattttatta	caacataaat	gtactttttc	2220
tctatacttg	ggaccaatat	tgaaattht	gattttatta	caccaaatt	taaattttat	2280
tacattaata	tttaaaattg	tattagaggt	ctcatgattt	ggtactacgg	gtctccgcat	2340
tatttccttt	ccaaatttcc	taatctgttt	caccaagggt	tctggacaac	tttagagacc	2400
ttttgtgaag	tttgaataaa	atctcttcga	gattttgata	attgcattag	ctttaggact	2460
taattggaat	agaattaaaa	tccttaaaac	aagctcttat	a		2501

<210> 3
 <211> 2251
 <212> DNA
 <213> Homo Sapiens

<400> 3

ggacccccgag	ccgccccccag	gtagccagga	ggggcctcag	cggcagccgc	aaactccagt	60
agccgcccgtg	gctgcccgtg	gctggggcgg	agggcagcca	gagctgggga	ccaaggtccc	120
gcgccacctg	cgcgcacagc	ctcacacctg	aacgctgtcc	tccgcgagac	gagaccggcg	180
ggcactgcaa	agctgggact	cgtctttgaa	ggaaaaaaa	tagcgagtaa	gaaatccagc	240
accattcttc	actgacccat	cccgtgcac	ctcttgtttc	ccaagttttt	gaaagctggc	300
aactctgacc	tcgggtgtcca	aaaatcgaca	gccactgaga	ccggctttga	gaagccgaag	360
attttggcagt	ttccagactg	agcaggacaa	ggtagaagca	ggttggaggc	gggtccagga	420
catctgaggg	ctgacccctg	gggctcgtga	ggctgccacc	gctgctgccg	ctacaggtga	480
gatggcggtg	ggctgacgtt	ggggtcaacg	ggtagagaac	gcagggatgc	ggccctcgcc	540
gaagagagcc	aagaagggaa	gagcgcgctc	tccaaattgc	ttttgtaact	tgttttcagt	600
gagcatttta	ttgattcaga	atctatcgag	aatagcacta	gagagctact	tttcccttga	660
gatgggtctt	attcatcttg	gcaatggagt	gagttggatt	gtggggagga	agagggaatgg	720
gaaaaatcagt	ttataaatat	taatgtcagc	aagagtgtgc	tgttggcagg	acgtatcgcg	780
agcctggaga	ttttgggtggc	cgcagttggt	aagtggctac	aatccagaaa	gtaggatcga	840
gttgctcccc	ttgtcttate	agtgtatcgt	ttctcgggcg	cggttctaac	accttacaag	900
tggtaatthc	cgtcacgggc	agctttgtct	ctcttctacc	atccccagac	ccagccttgc	960
actccaaggc	tgcgcacccg	cagccactat	catgtccact	ccgggggtca	attcgtccgc	1020
ctccttgagc	cccgaaccgg	tgaacagccc	agtgaccatc	ccggcggtga	tgttcactct	1080
cggggtgggtg	ggcaacctgg	tggccatcgt	ggtgctgtgc	aagtgcgcga	aggagcagaa	1140
ggagacgacc	ttctacacgc	tggtatgtgg	gctggctgtc	accgacctgt	tgggcacttt	1200
gttggtgagc	ccggtgacca	tcgccacgta	catgaagggc	caatggcccc	ggggccagcc	1260
gctgtgcgag	tacagcacct	tcattctgct	cttcttcagc	ctgtccggcc	tcagcatcat	1320
ctgcgccatg	agtgtcgagc	gctacctggc	catcaacgat	gcctatttct	acagccacta	1380
cgtggacaag	cgattggcgg	gcctcacgct	ctttgcagtc	tatgcgtcca	acgtgtctct	1440
ttgcgcgctg	cccaacatgg	gtctcggtag	ctcgcggtcg	cagtaccag	acacctggtg	1500
cttcacgcac	tggaccacca	acgtgacggc	gcacgcccgc	tactcctaca	tgtacgcggg	1560
cttcagctcc	ttcctcatte	tcgccaccgt	cctctgcaac	gtgcttgtgt	gcggcgcgct	1620

gctccgcatg	caccgccagt	tcattgcgccc	cacctcgctg	ggcaccgagc	agcaccacgc	1680
ggccgcggcc	gcctcggttg	cctcccgggg	ccaccccgt	gcctccccag	ccttgccgcg	1740
cctcagcgac	tttcggcgcc	gccggagctt	ccgccgcatc	gcggggcgccg	agatccagat	1800
ggtcatctta	ctcattgcca	cctccctggt	ggtgctcatc	tgctccatcc	cgctcgtggt	1860
gagtgaccgg	ggctggggcc	ctactcgccc	tttttctcgc	atccacctcc	cgctcgtggt	1920
ccccgctccc	tgctttccct	ctgagtcctt	ggcagtgaac	gtgtcgccct	taggtcgggg	1980
ctgggattcc	cacactggtt	ctcagagcag	gcccacccct	ctttgaagtc	ccaaccctaa	2040
cgagatttag	caggtgcttt	gcccctacat	ccccagttt	atgttcccgg	aagcctgggt	2100
ttctttctcc	accgagacag	cccttaccct	ttgctgcctg	acactggccg	agtcttccaa	2160
gaaaaccccc	cgccccctct	gttagacgtg	gaggggagcc	tgctgtagtg	tgacttagcc	2220
cattcctccg	tactgtgaac	tgtgaactgc	a			2251

<210> 4
 <211> 2586
 <212> DNA
 <213> Homo Sapiens

<400> 4

ccccgcgggg	cgccgggagta	gccccgctgg	gcgctcgccag	ccgcggggagt	caagccccct	60
ccccaggtgc	aggcataaaa	gtttatggct	cttgaacaat	gcggggcaga	ggtttttcca	120
agcaacgtct	aattggccgc	ttctaattaa	ggaaagagag	gcttccagct	ctatggcaac	180
ccaagcaggg	cagcttcagg	ctaaagggtac	tttagaataa	taagatcatt	ctaagaaatg	240
gaatgtctca	ctggacaccc	gaacagggttc	tctgtcattg	gaattggtgt	gtactgtact	300
tcaaccagta	ctcttgtgtg	gagggaggcg	accagtccta	ggaaagtcaa	ctacagaaag	360
aggtgacctc	cgaaaggatt	gtcttagcgc	tattagaata	catgtgacca	caccaaagc	420
ccaggcggac	accgcagcc	agctcggatt	tggacaattc	aacattgctg	gcagaactga	480
agggacaag	ttaccccaac	cccatccctt	gtacgcgtag	tgctgagtga	gttgggggtg	540
ggaggacagc	ggttcgttta	ttgccccctt	ttaaaatctg	agatctgaaa	atatggaggt	600
cccattcggt	ttcccagctc	ttgattgcca	acaaaaaac	aaatcccgt	ggctacattt	660
tctcctcatt	ccaaaatagc	aaccttatgg	cttgtattaa	gcccctcaga	agtttatctc	720
atattgctctg	ggccagggag	ggaacaatgc	taggaaaagt	caccggtgct	cttccatcct	780
cgcccccttc	aggggtcagg	atgtgcgggc	cgccgggcct	gtgatcccgg	aacgcttcc	840
gccatccctt	tgccggaact	tgaaggagct	gggagggtgt	gagagcagag	ttcagggctg	900
gtgcactctg	cggtgctgag	tgggcggcgc	gcccgggcgc	tcaggccggg	ggacctgtag	960
tcgccttacc	cgggagggga	aaatacgtag	ctggaggggc	tgccgcgtgc	gggttggtgat	1020
ccgttaccct	atcggtcacc	ctggggtctc	cccaagcctc	taggtagggc	tgtgagagtc	1080
ccctagagct	gaagcccccg	aggctgacct	gtgggtcttg	ctgctatggg	aacccgggtg	1140
gtccaaagaa	gcctttcttc	cgggcacctg	gaattccagt	ttagtgtggg	gcatcgggga	1200
agtggcgctg	gggggctggg	ttgggggacc	tcagccggca	gctccggaga	gggcctacc	1260
ttggggctgc	tgggtgaggc	cggcacgatt	cttggtccca	aaaggaaagt	ttctgcttct	1320
tggttctggc	cgagaagcca	aagacttatt	ttgagagcgg	agagagaaat	gttattggta	1380
acgttttctt	tggaaagtcc	gagaggggtc	ttctggacac	actacctagt	gcccccaaac	1440
cagagaagta	gtttttcttt	ggtgcctggg	ctcagaagtc	gccactcact	cagcccatgg	1500
ttcgaaatca	gcattgggaag	cgccggggca	aggctctgtc	ggagactaga	ggcctgacct	1560
tcgggaggag	cccctggggg	atgggggacc	cattctcctg	cttgctctgg	ttcccacctg	1620
ggacgcctcc	gtaggagccc	agaaagacga	tccactacat	ggtcccggga	cagagcagcg	1680
cgcccaactt	tgagggaact	ttgtgcgcct	ctctgaggcc	ctagctttcc	aaggcaccgc	1740
cgctcgttct	tctttcccta	gaccgaaact	ggggaagagt	gtgggcgctt	ctttgccccg	1800
atgagttcgc	ctcccaaac	gcctactctg	gctgcaccag	agcatctggg	aaactctgaa	1860
aggtgcccag	gcctcacaca	gcagcgtctc	ccactcagc	ctctgtcttt	gggttttttc	1920
aagagagctc	ctacctcatg	cctcgggtctt	tcttcgatgt	cggttccccg	aggtaggcac	1980
ggagtccctc	tgaagcagtc	tgccatctct	tgcctcttgc	gtgtaaagtt	agagtttact	2040
ttgttggggg	aaggggaggt	agaaaagatc	acagttggga	aagtgcgctt	ttcgcttctg	2100
tctataaaag	tgctcaaga	ctgtcatcgc	gattgttagg	agagctatca	acgtctaggg	2160
gtgggggtct	cttggggctg	ccctcgccc	ttcccaaac	cccaggttcc	taaaacccta	2220
ccctttatta	atatttcacg	ggattcaggc	tggcaccgct	gggaggacct	cgccctagcat	2280
atgttcccaa	actctggcta	aaggcaggct	cctgccttct	ctggagcctc	ttttctcgga	2340
ggaagagacg	cgtcaactcc	actcactccc	ctgtgagcca	tcctagggct	ctgtgccccg	2400
tgacccccct	gacggccctt	cggggtctgc	gcgcagtcct	tagccgcaaa	gtgctgcaag	2460
ctttattttt	aaaagaacaa	tccgaccgaa	gagctcgggg	accaaagaga	aaaaaaataa	2520
tgggggt	gtcatcactg	cggcgatact	gtggcgagg	actttggcga		2580

<210> 5
 <211> 2501
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 5

ttgttgata	gaatatttta	ttatttaggt	attatgtcga	gtatttaata	gttttttttt	60
ttgttttttt	tttttttttt	atattgtatt	ttggagttaa	ttatagtgtt	tgttgttttt	120
ttgtttgtgt	tataagtttt	tattatttag	ttttatttta	taagtgaaga	tatttagtat	180
ttggattttt	gtttttgtat	tagtttgtaa	aggataatag	tttttagttt	tatttatgtt	240
tttataaaa	atatgattta	gtttttttta	atggttgat	taaatagaat	tttaaagata	300
taataataa	attaattttt	tttttattat	aaaaattttt	tgttgaattt	gatttatatt	360
aaattaacga	gttttgtttt	atgaaagatt	ttttggataa	atttgatagt	tgatggaata	420
ggagaagttg	tttggtatgt	ttaaagttta	taagagatta	atatttagaa	taaataggaga	480
tttgtaaaat	aatagaaagt	aggtagtaaa	gttaaagaaa	atagtttaag	gtatagttat	540
taaaaggaac	gtgattatgt	tttttgtagg	gatatgggtg	gagttggaag	tcgttagttt	600
tagtaaaatt	atataggaat	agaaaattag	cgagatcgta	tggtttttat	tataagtggg	660
agttgaataa	tgagaatata	tggttatatg	gcggcgatta	atataattg	gtgtttgttg	720
agcggggtgt	tggggagggg	gagttatagg	aagaatagtt	aagggatatt	gggtttaata	780
tttggtgat	gggatgattt	gtatagtaaa	ttattatggc	gtatataatt	atgtaataaa	840
tttgatatatt	ttttatatgt	attttagaat	tttaaataaa	agttggacgg	ttaggcgtgg	900
tggtttacgt	ttgtaatttt	agtatttttg	gaagtcgagg	cgtgtagatt	atttaaggtt	960
aggagttcga	gattagttcg	gttaatatgg	tgaaatttcg	tttttattaa	aaatataaaa	1020
attagttaga	tggtgtacgt	atttataatt	ttatttatcc	gggaggttga	agtagaattg	1080
tttgaattcg	agaggcggag	gttgtagtga	gtcgtcgaga	tcgcgttatt	gtatttttagt	1140
ttgggttata	gcgtgagatt	acgttatata	ataaaataaa	ataatataaa	ataaaataaa	1200
ataaaataaa	ataaaataaa	ataaaataaa	ataaaataaa	ataaaaaaat	aaaaataaat	1260
aaaaataaaa	aaagtaattt	tttttttttt	aagcggtttt	tatttttttt	ttttgttttg	1320
tgaagcgggt	gtgtaagttt	cgggatcgta	gcggttttag	ggaatttttt	ttcgcgatgt	1380
ttcggcgct	tagttcggtt	cgtatatttc	gttcgggttt	ttttttgttt	gtttgtttat	1440
tttttaggtt	tcgttgggga	tttgggaaag	agggaaaggt	tttttcgggt	agttgcgcgg	1500
cgatttcggg	gatttttaggg	cgtttttttg	cggtcgacgt	tcggggtgta	gcggtcgctc	1560
gggttcgggt	cgccgggagt	tcgcgggatt	ttttagaaga	gcggtcggcg	tcgtgattta	1620
gtattggggc	ggagcggggc	gggattattt	ttataagggt	cggaggtcgc	gagggtttcg	1680
ttggagtttc	gtcgtcgtag	ttttcgttat	tagtgagtac	gcgcggttcg	cgttttcggg	1740
gatgggggtt	agagttttta	gtatgggggt	aattcgtagt	attaggttcg	ggttttcggg	1800
aggggttttc	gtttatttcg	agattcggga	cgggggttta	ggggatttag	gacgttttta	1860
gtgtcggttag	cggttttttag	gggggttcgga	gcgtttcggg	gagggatggg	atttcggggg	1920
cggggagggg	gggtagattg	cgtttatcgc	gttttggtat	tttttttcgg	gttttagtaa	1980
attttttttt	gttcgttgta	gtgtcggttt	atatacgtgt	ttatttttta	gttcgaggta	2040
ggagtatgtg	tttggtaggg	aaggagggta	gggggttggg	ttgtagttaa	tagtttttcg	2100
tttattcggg	gagattcgaa	tttttttatt	ttttcgtcgt	gtggttttta	tttcgggttt	2160
tttttttggt	tttcgttttt	ttcgttatgt	ttgtttttcg	ttttagtgtt	gtgtgaaatt	2220
ttcggaggaa	ttgttttttt	tggttttttt	ttgtattttt	gatttttttt	cgggttggtg	2280
cgaggcggag	tcgggttcggt	ttttatatatt	cgtatttttt	ttttttcgta	ggtcggttcg	2340
cggttttcgc	tatgttgttg	gtagattagg	gttagagttg	gaaggaggag	gtggtgatcg	2400
tggagacgtg	gtaggaggtt	ttatttaaag	ttttttgcgt	aagtgattat	gttcgggtaa	2460
ggggaggggg	tggtgggttt	tagggggttg	tgattaggat	t		2501

<210> 6

<211> 2501

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 6

gatttttagtt	atagtttttt	aaggtttagt	atattttttt	tttgttcggg	tatggttatt	60
tacgtaggag	gttttgagt	agtttttttg	ttacgttttt	acggttatta	tttttttttt	120
ttagtttttg	ttttgatttg	ttagtagtat	gcgtaggggt	gcgtagcggg	ttgcggggag	180
ggagaagtac	gagatgtggg	gatcgggtcg	atttcgtttc	gtagtaattc	ggggaggggt	240
taggagtgtg	gggagggagt	agggaaatag	gttttttcga	agattttata	taataattgg	300
gcgggagtag	ggatagggcg	gagagggcgg	gaataggaa	gaggttcggg	gtaaaagtta	360
tacgagggag	ggataagggg	gttcggattt	tttcgggttg	gcgaggggtt	gtgggttgta	420
gttttagttt	ttgttttttt	ttttgttag	atataatgtt	ttatttcgaa	ttgggaaata	480
gattacggtg	tagggcggta	ttgtagcgaa	taaagaaaag	tttggtggag	ttcgggggag	540
gatgttaagg	cgcggtgagc	gtagtttgtt	tttttttttc	gttttcgggg	ttttattttt	600
tttcgagggc	tttcgggttt	tttgaaagtc	gttaacggta	ttggggacgt	tttgggtttt	660
ttaggttttc	gttatggggt	tcgaggtggg	cgaggagttt	tgctgggagt	tcgggtttga	720
tggttcgggt	tggtttttat	ttgggagttt	tgagttttat	tttcggggac	gcgggtcgcg	780
cgtattttat	ggtggcgaag	attgcggcgg	cgaaatttta	gcgaagggtt	cgcgggtttc	840
gagttttata	aggggtgggt	cgtttcgttt	cgtttttagt	ttgagttacg	gcgtcgggtc	900

tttttttggg	gggttttcg	gatttttcgt	ggttttagtt	tggcggtcg	ttgtatttcg	960
ggcggtcggt	gtagaggggc	gttttggagt	tttcggagtc	gtcgcgtagt	tggtcgggga	1020
agtttttttt	gttttttag	gttttttagc	gggttttagg	agtaaataga	tagtaggaa	1080
aggatcgtag	cgaagtgtgc	gtagcgaatt	ggcgcgctcg	gatatcgcg	ggggaaattt	1140
tttaagatcg	ttgcgatttc	ggagtttgta	tattcgtttt	atagggtagg	ggagaggggt	1200
ggaggtcggt	tagaggaaa	gaaattgttt	tattttattt	tattttattt	tattttttta	1260
ttttatttta	ttttatttta	ttttatttta	ttttatttta	ttttatttta	ttttgtgtta	1320
ttttatttta	ttttatgacg	tagttttacg	ttgtggttta	ggttggagtg	tagtggcgcg	1380
atttcggcgg	tttattgtaa	ttttcgtttt	tcgggtttta	gtaattttgt	tttagttttt	1440
cgagttagtg	gaattatagg	tcggtgttat	atttggttga	ttttgtatt	tttagtagag	1500
acgggggttt	attatgttgg	tcgggttggt	ttcgaatttt	tgatttttag	tgattttgtac	1560
gtttcgggtt	tttaaagtgt	tgggattata	ggcggtgagt	attacgtttg	gtcggttaat	1620
ttttatttga	agttttgggg	tatatgtaga	ggatgtgtag	gtttgttata	taggtgtgtg	1680
cgttatgatg	gtttgttgta	tagattattt	tattatttag	gtattaaagt	tagtattttt	1740
tagttatttt	ttttgggtatt	tttttttttt	agtatttcgt	ttaataggta	ttagtgtgtg	1800
ttgatcgctg	ttatgtgatt	atgtgttttt	attgttttag	ttttatttat	aagtgtgatt	1860
atgcgggttt	gttgggtttt	tgtttttgtg	tgagtttggt	gaggttaacg	gttttttagt	1920
ttatttatgt	ttttgtaaa	gatattgatta	cgtttttttt	agtggttgtg	tttttaggtta	1980
tttttttttg	ttttgttgtt	tattttttgt	tgattttagt	attttttatt	atttttagata	2040
ttgatatttt	gtttgtttta	gatattgata	atagtttttt	ttattttatt	aattgttaag	2100
tttgtttaag	gagtttttta	tgaataaaaa	ttcgttaatt	taagtgtaat	taaatttagt	2160
aagggtattt	tgtgggtggg	aagaggttgg	tgtttatgtt	gtatttttaa	aattttattt	2220
aatgtagtta	ttaaaaagaa	ttagattatg	ttttttgtgg	gaatatggat	ggagtttagag	2280
gttattattt	ttagtaaatt	aatgtaggaa	tagaaattta	aatattggat	gtttttattt	2340
gtaagtggga	gtttaaagt	gagaatttat	aatataaata	aggaaataat	agatattgtg	2400
gttgatttta	gggtgttaga	tgggaggaag	gagaggagta	gaaaagagaa	ttattgggta	2460
ttcgggtata	tatttgggtg	atgaaatatt	ttgtataata	a		2501

<210> 7

<211> 2501

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 7

tttgtaaattg	gagatatttt	tattattttt	atagtattat	atgtttttta	agtttgtatt	60
tatatatttg	gtgataaatt	aaggataaga	ttttttttta	tttttgtgag	gatgattata	120
gtatgattgg	atgggtttgt	tatgattttt	attttttttt	gtgtttttat	tatcgtttta	180
tttaattttag	ttttttttta	tagggtagta	tagaatttaa	ttagtagaaa	gagatttagt	240
tatgtagatt	agagatttgt	ttaagtgcg	gtatgtaaga	attaggaagg	aaagtttttt	300
gtttaaatat	taataggttt	tttttttaaa	gtaattatta	ttttttaaat	tttaattata	360
agggtgatagt	atttttaaat	taattaaatt	agaatttcgg	gttggaata	tttaaatatg	420
attttattagt	attttttatt	aattattggg	tttttagggt	tttaagttta	tttatttaga	480
attttatttt	taatattatt	ttattaattt	tagttgtaaa	taagagaata	tttaaagggt	540
gaggaatttt	tagcggtaaa	gttttgttta	cgttaagtaa	taaaggataa	gttagttttt	600
gttgtgatta	ttttgttgta	ttgataaggt	acgtattttt	atttaaggat	ttaaattttt	660
atttttttta	agaattgggt	taaaatcgat	aaattaaatt	tatttacggt	ttattgatta	720
aaggttgttg	tataataagt	ttttgttatg	tttagtagtt	ggatttatag	cgttagaaat	780
ttataattgt	ttgatttttt	tttttattat	attgcgaaaa	ttgtttttta	aatgtaatta	840
attttaaaat	tttaatagta	tcgtggtttag	gcgtgggtgg	ttattattgt	aatattaata	900
ttaggtatag	gcgaggggat	tgaggtttag	atatcgaaat	tagtttgagg	aatatacgga	960
gattcgggtt	ttggaaaaat	aattagtttt	gcgtgggtgg	gggcgcgagg	tttcgggtta	1020
tcgggagggt	atgatgattt	atgatgatat	tgtattatag	tttgcgcgac	ggtttatggt	1080
agtaagtttt	ggagtatttg	aaataagttg	tggtgggtat	tttatttatt	ggagagcgat	1140
tagtgattga	tgtttattta	tagcgttagg	agacgtatgt	ttcgatagta	gtataaattt	1200
agtaggcgcg	aataaatggt	aaagagaaa	tgggtaaata	agtattacgg	tttttttagt	1260
gagaaagtgg	gggtttttaa	aagggttttt	tggtgataga	aagggtacgt	taattatcga	1320
aatcgtagag	ggtgcgggtt	tggcgtttga	gcgcgttagt	tatatattatg	gcggtgatcg	1380
tttcgtggtt	ggcgtgtttt	gtataggtta	cgcggtttcg	gattacgttt	tttaggaata	1440
tttttagtat	ttcgcgagtt	ttttcgtaga	tgaggtcgga	gatgcgtttt	acgtcgtcgc	1500
ggcgagtaag	gcgtcggatg	gtcggtttgg	tgatgttttg	gatattgtcg	cgtagtattt	1560
tacgggtggc	tttagcgtcg	ttttgtttta	gatttttttc	gtttttgtcg	cggtagata	1620
tgacgagtaa	gaggagtttt	atttaacgtt	tttgtgaggt	tttggtttga	ggttagcgtt	1680
ttatacgata	gttggcggat	cgaattgaga	atttgaaaga	agtcggcggg	aagtttcgtt	1740
tcggtggggg	aggggaaatt	taaaagggtta	aatcgaaata	gggggaaaaa	aaaagcgagt	1800
tttttgtttt	cgtgttttga	attttgtaac	gtgtatagta	ttttgttatt	acgttatgag	1860
gttttaaaaa	attgtttttg	aacgtagaag	atatatatta	atattgtggg	aaatataaga	1920
aaggataaga	aattaagaaa	ttataatgtt	attttattat	ataggttagt	taattatgta	1980

ttttgtagag	tagttgtata	tattttttta	agaaaatgta	tatagtgttg	tatatggagt	2040
tttgtaattt	ttttatat	attataattt	aattaatttt	tattaaagag	ataaaagtga	2100
tgttttgggt	ttttatgttt	ttaggaatta	ttaatagtta	taattagttt	tttagtaatt	2160
ttttaacogg	ttgtatttta	aaaataatgt	tttttatatt	taatataaat	gtattttttt	2220
tttatatttg	ggattaatat	tgaaatttat	gattttatta	tattaaaatt	taaattttat	2280
tatattaata	tttaaaattg	tattagaggt	tttatgattt	ggtattacgg	gttttcgtat	2340
tatttttttt	ttaaattttt	taatttggtt	tattaaggtt	tttggaataa	tttagagatt	2400
ttttgtgaag	tttgaataaa	attttttcga	gatttttgata	attgtattag	ttttaggatt	2460
taattggaat	agaattaaaa	tttttaaaat	aagttttttat	a		2501

<210> 8

<211> 2501

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 8

tataagagtt	tgttttaagg	attttaattt	tattttaatt	aagtttttaa	gttaatgtaa	60
ttattaaaat	ttcgaagaga	ttttatttaa	atttttataa	aggtttttaa	agttgttttag	120
aaatttttgg	gaaatagatt	aggaaatttg	gaaaggaaat	aatgcggaga	ttcgtagtag	180
taaattatga	gattttttaa	ataattttta	atattaatgt	aataaaaatt	aaatttttgg	240
gtaataaaaa	tataaatttt	aataattgggt	ttaagtatat	agaaaaagta	tatttatggt	300
gaatgtggaa	aataattttt	ttaaaatata	gtcgattaaa	aaattgttgg	ggaattgatt	360
ataattattg	ataattttta	agaaatatag	atattaaaat	attattttta	tttttttaat	420
agaaattggg	taaattataa	ttaatataga	gaggttataa	aattttatat	ataatattgt	480
atataatttt	ttggaaaaat	atgtgtaatt	gtttttgtaa	atataatgatt	aattagtttg	540
tgtgatggga	taataattgt	gttttttaat	tttttgtttt	tttttgattt	ttttatagta	600
ttgatgtata	ttttttgcgt	ttaaaagtaa	ttttttaaag	ttttataacg	tggtataaaa	660
atattatgta	cgttataaaa	tttagaatac	ggaaataaga	agttcgtttt	tttttttttt	720
ttatttcggg	ttggtttttt	agattttttt	ttttttatcg	ggcggggatt	tttcgtcgat	780
tttttttagg	tttttagttc	ggttcgttaa	ttgtcgtata	aaggcgttgt	tttaggttag	840
agtttttata	aagcgttggg	tgagattttt	ttgttcgttt	atgtttgggt	gcggtaagg	900
cgggaagggt	tttggtaaaag	gcggcggtta	gcgttatcgt	aaagtattgc	gcgataatat	960
tttaggtatt	attaagtcgg	ttattcggcg	ttttgttcgt	cgcgccggcg	tgaagcgtat	1020
tttcgggttt	atttacgagg	agattcgcgg	gggtgtgaag	gtgttttttg	agaacgtgat	1080
tccgggacgt	gtgatttata	tagagtacgt	taagcgttaag	acgggttatc	ttatggatgt	1140
ggtttacgcg	tttaagcgtt	agggtcgtat	tttttacggg	ttcgggtggt	gagcgttttt	1200
ttttattaat	aaaagggttt	tttttaggggt	ttttattttt	tagttgagga	gtcgtgatgt	1260
ttgtttgttt	agtttttttt	tattattttg	tcgcgtttgt	tgagtttgtg	ttgttatcgg	1320
agtatgcgtt	tttagtcggt	gtaagtaggt	attagttatt	aatcgttttt	tagtaataaa	1380
aatattttaat	ataatttggt	ttagggtgtt	tagagtttat	tgatatgggt	cgtcgcgtag	1440
attgtagtgt	agtgttatta	tggtttattg	tagtttttcg	attagtcgga	atttcgcggt	1500
tattattacg	taaggttaat	tattttttta	aagatcgggt	tttcgtgtgt	tttttaggtt	1560
agtttcgata	ttttgggttt	aatttttttc	tttatgttta	atgttggtat	tatagtagtg	1620
agttattacg	tttggttacg	atattgttga	ggtttttagg	ttagttatat	ttaaggggta	1680
attttcgtag	tgtagtgggg	aggaaagtta	agtagttata	ggtttttggc	gttgtgaatt	1740
taattgttga	atatagtaag	aatttattat	gtaataattt	ttaattagtg	gatcgtaaat	1800
aagtttagtt	tatcgggttt	ggtttaattt	ttgagaaagg	tgagaatttg	aatttttgag	1860
tagaaataac	tagttttatta	gtataataaa	gtgattataa	taaagattaa	tttatttttt	1920
gttattttaac	gtgggttagag	ttttatcgtt	gagaattttt	tagtttttga	gtgttttttt	1980
attttataat	gaagttgata	agatggtatt	aaaagtgaga	tttttagtaa	gtaaatttta	2040
aggtttgaag	gattagtgat	taatgggaag	tgtaataaag	ttatatttga	ggttatttaa	2100
ttcgagattt	tgattttaatt	ggtttaagga	tattattatt	ttgtgggtta	gatttgaaaa	2160
ataataaatt	ttttaaggaa	ggaatttggt	ggatttttaa	taaaaaattt	ttttttttga	2220
tttttatatg	tcgttatttt	gataaatttt	tggtttatat	ggttggattt	ttttttgtta	2280
gttaaatttt	gtgttatttt	gtgaaaaaga	attgagatta	ataaaacggt	agtgagaata	2340
taggggaaaga	taaaaattat	agtaagttaa	tttagttatg	ttgtagttaa	ttttataagg	2400
ataggggaag	attttggttt	ttatttatta	tttaagtggt	gagtataaat	tttaaaaaata	2460
tatgatatta	taggaataat	gaagatgttt	ttatttgtaa	a		2501

<210> 9

<211> 2251

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 9

ggatttcgag	tcggttttag	gtagttagga	gcggttttag	cggtagtcgt	aaatttttagt	60
agtcgttcgt	gttggttcgtg	gttggggcgg	agggtagtta	gagttgggga	ttaaggtttc	120
cgtttatttg	cgcgtatagt	tttataattg	aacggttggtt	tttcgtagac	gagatcggcg	180
ggtattgtaa	agttgggatt	cgtttttgaa	ggaaaaaaa	tagcgagtaa	gaaatttagt	240
attatttttt	attgatttat	ttcgttgat	tttttggttt	ttaagttttt	gaaagtgggt	300
aattttgatt	tcggtgttta	aaaatcgata	gttattgaga	tcggttttga	gaagtcgaag	360
atttggtagt	ttttagattg	agtaggataa	ggtgaaagta	ggttggaggc	gggttttagga	420
tatttgaggg	ttgattttgg	gggttcgtga	ggttggtatc	gttggtgtcg	ttatagggtga	480
gatggcgttg	ggttgacgtt	ggggttaacg	ggtagagaac	gtagggatgc	ggttttcgtc	540
gaagagagtt	aagaagggaa	gagcgcgttt	tttaaatgt	ttttgtaatt	tgtttttagt	600
gagtatttta	ttgatttaga	atztatcgag	aatagtatta	gcgagttatt	tttttttga	660
gatgggtttt	atztattttg	gtaatggagt	gagttggatt	gtggggagga	agaggaatgg	720
gaaaaattag	ttataaatat	taatgttagt	aagagtgtgt	tggtggtagg	acgtatcgcg	780
agtttgagga	ttttgggtgg	cgtagttggg	aagtgggtat	aatttagaaa	gtaggatcga	840
gttggttttt	ttgttttatt	agtgtatcgt	ttttcgggcg	cgggtttaat	attttataag	900
tggtaatttt	cgtttacggg	agttttgttt	tttttttatt	attttttagat	ttagttttgt	960
attttaaggt	tgcgtatcgt	tagttattat	tatgtttatt	ttcgggggta	attcgttcgt	1020
ttttttgagt	ttcgatcggg	tgaatagttt	agtgattatt	tcggcgggtga	tgttttttt	1080
cgggggtggg	ggtaatttgg	tggttatcgt	ggtgtgtgtg	aagtcgcgta	aggagtagaa	1140
ggagacgatt	ttttatacgt	tggtatgtgg	gttggtgtgt	atcgatttgt	tggttatttt	1200
gttggtgagt	tcggtgatta	tcgttacgta	tatgaagggt	taatgggtcg	gggggttagtc	1260
gttggtgcgag	tatagtattt	ttattttgtt	tttttttagt	ttgttcgggt	ttagtattat	1320
ttgcgttatg	agtgtcgagc	gttatttggg	tattaatatt	gtttattttt	atagttatta	1380
cgtggataag	cgattggcgg	gttttacgtt	ttttgtagtt	tatgcgttta	acgtgttttt	1440
ttgcgcgttg	tttaatatgg	gtttcggtag	ttcgcgggtg	tagtatttag	atatttggtg	1500
ttttatcgat	tggtattatta	acgtgacggc	gtacgtcgtt	tatttttata	tgtagcgggg	1560
tttttagtttt	ttttttattt	tcgttatcgt	tttttgtaac	gtgtttgtgt	gcggcgcgtt	1620
gtttcgtatg	tatcgttagt	ttatgcgtcg	tatttcgttg	ggtatcgagt	agtattacgc	1680
ggtcgcggtc	gtttcgggtg	tttttcgggg	ttatttcgtt	gttttttttag	ttttgtcgcg	1740
tttttagcgat	tttcggcgctc	gtcggagttt	tcgtcgtatc	gcgggcgtcg	agatttagat	1800
ggttattttta	tttattgtta	tttttttggg	ggtgtttatt	tgttttattt	cgttcgtggg	1860
gagtgatcgg	ggttgggggt	ttatttcgggt	tttttttcgt	atttattttt	cgcgtttatt	1920
tttcggttttt	tgtttttttt	ttgagttttt	ggtagtgaac	gtgtcgtttt	taggtcgggg	1980
ttgggatttt	tatatgtttt	tttagagtag	gtttaatttt	ttttgaagtt	ttaattttaa	2040
cgagattttag	taggtgtttt	gtttttatat	tttttagttt	atgttttcgg	aagtttgggt	2100
tttttttttt	atcgagatag	ttttttattt	ttgttggttg	atattgggtcg	agtttttttaa	2160
gaaaattttt	cgtttttttt	gttagacgtg	gaggggagtt	tggtgtagtg	tgatttagtt	2220
tatttttttcg	tattgtgaat	tgtgaattgt	a			2251

<210> 10

<211> 2251

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 10

tgtagtttat	agtttatagt	acggaggaat	gggttaagtt	atattatagt	aggttttttt	60
ttacgtttta	tagagggggc	gggggggttt	tttggaagat	tcggttagtg	ttaggtagta	120
aggggtaagg	gttggttcgg	tggagaaaaga	aatttagggt	ttcgggaata	taaattgggg	180
gatgtagggg	taaagtattt	gttaaatttc	gttaggggtg	ggatttttaa	gagggttggg	240
tttggtttga	gaaatagtgt	gggaatttta	gtttcgattt	aaaggcgata	cgtttattgt	300
taaggattta	gagggaaagt	agggagcggg	gaatggacgc	gggaggtgga	tgcgagaaaa	360
aggtcgagta	gggttttagt	ttcggttatt	tattacgagc	gggatggagt	agatgagtat	420
tattagggag	gtggtaatga	gtaagatgat	tatttggttg	tcggcggttcg	cgatgcggcg	480
gaagtttcgg	cggcgctcgaa	agtcggttag	gcgcggtaag	gttggggagg	tagcgggggtg	540
gtttcggggg	gtaatcgagg	cggtcgcggg	cgcgtggtgt	tggtcgggtg	ttagcgaggt	600
gcggcgtag	aattggcggg	gtatgcggag	tagcgcgtcg	tatataagta	cgttgtagag	660
gacggtggcg	agaatgagga	aggagttgaa	gttcgcgtat	atgtaggagt	aggcggcgtg	720
cgtcggttacg	ttggtgggtt	agtcgatgaa	gtattaggtg	tttgggtatt	gtagtcgcga	780
gttatcgaga	tttatgttgg	gtagcgcgta	aaagagtacg	ttggacgtat	agattgtaaa	840
gagcgtgagg	ttcggttaatc	gtttgtttac	gtagtgggtg	tagaaatagg	tatggttgat	900
ggtaggtag	cggttcgatat	ttatggcgta	gatgatgttg	aggtcggata	ggttgaagaa	960
gagtagaatg	aaggtgttgt	attcgtatag	cgggtgggtt	tcgggttatt	ggttttttat	1020
gtacgtggcg	atgggttatcg	ggtttattaa	taaagtgttt	aatagggtcg	tgatagttag	1080
tttataatatt	agcgtgtaga	aggtcgtttt	ttttgttttt	ttgcgcgatt	tgtatagtat	1140
tacgatgggt	attaggttgt	ttattatttc	gaagatgaat	attatcgtcg	ggatgggtat	1200

tgggttggtt	agtcggtcgg	ggtttaagga	ggcggacgaa	ttgatttcgg	gagtggatat	1260
gatatgtggt	ggcgggtgcgt	agttttggag	tgtaagggtt	ggtttgggga	tggtagaaga	1320
gagataaagt	tgctcgtgagc	ggaaattatt	atattgtaagg	tgtttagattc	gcgttcgaga	1380
aacgatatat	tgataagata	aggggagtaa	ttcgattttta	ttttttggat	tgtagttatt	1440
tattaattgc	ggttattataa	attttttaggt	tcgcgatacg	ttttgttaat	agtataattt	1500
tgttgatatt	aataattata	aattgatttt	tttatttttt	ttttttttta	taatttaatt	1560
tattttattg	ttaagatgaa	taagattttat	tttaagggaa	aagtagttcg	ttagtgttat	1620
tttcgataga	ttttgaatta	ataaaatggt	tattgaaaat	aagttataaa	agtaatttgg	1680
agagcgcgtt	tttttttttt	tggttttttt	cggcgagggt	cgtatttttg	cgttttttat	1740
tcgttgattt	taacggttagt	ttaacggtat	tttatttgta	gcggtagtag	cgggtgtagt	1800
tttacgagtt	tttagggtta	gttttttagat	gttttggtt	cgtttttaat	ttgtttttat	1860
tttggtttgt	ttagtgttga	aattgttaaa	ttttcggttt	tttaaagtcg	gttttagtgg	1920
ttgtcgtatt	ttggatatcg	agggttagagt	tgtagttttt	taaaaatttg	ggaaataaga	1980
gggtgtagcgg	gatgggttag	tgaagaatgg	tgtaggtttt	tttattcgtt	attttttttt	2040
ttttaaaagac	gagtttttagt	tttgtagtgt	tcgtcggttt	cgtttgcggg	aggatagcgt	2100
ttaggtgtga	gggtgtgcgc	gtaggtggcg	cggagttttg	gttttttagt	ttggttggtt	2160
ttcgttttag	ttacgggtag	tacgggcggt	tattggagtt	tcggttggtc	gttgaggtcg	2220
tttttggtta	tttggggggt	gttcgggggt	t			2251

<210> 11

<211> 2586

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 11

ttcgcgcggg	cgcgggagta	gtttcgttgg	gcgttcgtag	tcgcgggagt	taagtttttt	60
tttttaggtg	aggtataaaa	gtttatggtt	tttgaataat	gcggggtaga	ggttttttta	120
agtaacgttt	aattggtcgt	ttttaattaa	ggaaagagag	gttttttagt	ttatggtaaat	180
ttaagtaggg	tagttttagg	ttaaagggtat	tttagaataa	taagattatt	ttaagaaatg	240
gaatgtttta	ttggatatcc	gaatagggtt	tttggtattg	gaattgggtg	gtattgtatt	300
ttaattagta	ttttctgtg	gagggaggcg	atttagttta	ggaaagttaa	ttatagaaag	360
aggtagtttt	cgaagggtt	gttttagcgt	tattagaata	tatgtgatta	tattaaaagt	420
ttaggcggat	attcgtagtt	agttcggatt	tggataattt	aataattgtt	gtagaattga	480
agggaataag	ttattttaat	tttatttttt	gtacgcgtag	tgttgagtga	gttgggggtg	540
ggaggatagc	ggttcgttta	ttgttttttt	ttaaaatttg	agatttgaaa	atatggagggt	600
tttatttcgt	tttttagttt	ttgattgtta	ataaaaaaat	aaatttcgtt	ggttatatatt	660
tttttttatt	ttaaaatagt	aatttttatg	tttgtattaa	gtttttttaga	agttttattt	720
atttgttttg	ggtaggggag	ggaataatgt	taggaaaagt	tatcgggtgt	tttttatttt	780
cgtttttttt	agggtgtagg	atgtgcgggt	cggcgggttt	gtgatttcgg	aacgtttttt	840
gttatttttt	tcgcgcgaatt	tgaagggtt	gggagggtgt	gagagtagag	tttaggggtg	900
gtgtattttg	cgggtgttag	tgggcggcgc	gttcgggcgt	ttaggtcggg	ggatttgtag	960
tcgttttatc	cgggagggga	aaatacgtag	ttggaggcgc	tcggtcgtgc	gggttgtagt	1020
tcgttatttt	atcggttatt	ttgggggttt	tttaagtttt	taggtagggt	tgtgagaggt	1080
tttttagagtt	gaagtttcgg	agggtgattt	gtgggttttg	ttgttatggg	aattcgggtg	1140
gtttaaagaa	gttttttttt	cgggtatttg	gaatttttagt	ttagtgtggg	gtatcgggga	1200
agtggcgttg	gggggttggg	ttgggggatt	ttagtcggta	gtttcggaga	gggtttattt	1260
ttggggctgt	tgggtgaggt	cgttacgatt	tttggtttta	aaaggaaagt	ttttgttttt	1320
tggtttggcg	cgagaagtta	aagattttatt	ttgagagcgg	agagagaaat	gttattggta	1380
acgttttttt	tggaaagttc	gagaggggtt	ttttggatat	attatttagt	gtttttaaat	1440
tagagaagta	gttttttttt	ggtgtttggg	tttagaagtc	gttatttatt	tagtttatgg	1500
ttcgaataa	gtatgggaag	cgtcggggta	aggtttcgtc	ggagattaga	ggtttggttg	1560
tcgggaggag	tttttggggg	atggggattt	tatttttttg	tttggttttg	tttttatatt	1620
ggacgttttc	gtaggagttt	agaaagacga	tttattatat	ggtttcggga	tagagtagcg	1680
cgtttaattt	tgagggaatt	ttgtcgtttt	ttttgagggt	ttagtttttt	aaggtagcgt	1740
cgttcgtttt	ttttttttta	gatcgaaatt	ggggaagagt	gtgggcgttt	ttttgtttcg	1800
atgagttcgt	tttttttaac	gtttatttct	tttgtattag	agtatttggg	aaattttgaa	1860
agggtgtttg	gtttttatata	gtagcgtttt	ttttgttagt	ttttgttttt	gggttttttt	1920
aagagagttt	ttattttatg	tttcggtttt	ttttcgatgt	cgggttttcg	aggtaggtac	1980
ggagtttttt	tgaagtagt	tgttttattg	tggttttttg	gtgtaaagtt	agagttttatt	2040
ttgttggggg	aaggggaggt	agaaaagatt	atagttggga	aagtgcgttt	ttcgttttgt	2100
ttttaaaata	tgttttaaga	ttgttatcgc	gattgttagg	agagttatta	acgttttaggg	2160
gttgaagaag	aatttttgaa	ttttcggttt	tttttaaat	tttaggtttt	taaaatttta	2220
gtgggggttt	ttttgggggt	ggatttaggt	ttgtatcgtt	gggaggattt	cgttttagtat	2280
ttttttatta	atattttacg	aaggtaggtt	tttggttttt	ttggagtttt	ttttttcgga	2340
atgtttttta	attttgggtt	atttattttt	ttgtgagttt	ttttagggtt	ttgtgggttcg	2400
ggaagagacg	cgttaatttc	gcgggtttgc	gcgtagtttt	tagtcgtaaa	gtgttgtaag	2460
tgattttttt	gacggttttt	ttcgatcgaa	gagttcggga	attaaagaga	aaaaaaataa	2520

ttttattttt aaaagaataa gttattattg cggcgatatt gtggcggagg attttggcga	2580
tgggggt	2586

<210> 12
 <211> 2586
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 12

attttatcgt	taaagttttt	cgttatagta	tcgtcgtagt	gatgatttgt	ttttttgaaa	60
ataaagttat	tttttttttt	tttggttttc	gagtttttcg	gtcggaaagg	gtcgttaggg	120
gggttatttg	tagtattttg	cggttaagga	ttgcgcgtag	attcgcggag	ttgacgcgtt	180
ttttttcggg	ttatagagtt	ttaggatggt	ttatagggga	gtgagttagt	tagagtttgg	240
gaatatttcg	agaaaagagg	tttttagagaa	ggtaggagtt	tgttttcgtg	aaatattaat	300
aaagggatgt	taggcgaggt	tttttttagcg	gtgttagttt	gaattttagt	tttaagagat	360
ttttattagg	gttttaggaa	tttggggggt	tgggaagggt	cgaggggtta	gaaatttttt	420
tatagttttt	agacgttgat	agttttttta	ataatcgcg	tgatagtttt	gaggtatggt	480
ttaggaataa	ggcgaaaagc	gtattttttt	aattgtgatt	ttttttat	tttttttttt	540
taataaagta	aatttttaatt	ttatattaaa	ggggtataga	taggtaattg	tttttagagg	600
gatttcgtgt	ttatttcggg	gattcgatat	cgaagaaaga	tcgaggtatg	aggtagagat	660
ttttttgaaa	aaattttaag	atagaggttg	agttaggaga	cgttgttgtg	tgaggtttgg	720
gtatttttta	gagtttttta	gatgttttgg	tgtagtcgaa	gtaggcggtt	ggggaggcga	780
attttatcggg	gtaaagaagc	gttttatattt	tttttttagtt	tcgggttagg	gaaagaagaa	840
cggacggcgg	tgttttggaa	agttagggtt	ttagagaggc	gtataaagtt	tttttaaagt	900
tgggcgcggt	gttttgtttc	gggattatgt	agtggatcgt	ttttttgggt	ttttacggag	960
gcgttttagg	tgggaattag	agtaagtagg	agaatggggt	ttttattttt	taggggtttt	1020
tttcgatagg	taggttttta	gttttcgacg	aagttttgtt	tcggcggttt	ttatgttgat	1080
ttcgaattat	gggttgagtg	agtggcgatt	tttgagttta	ggtattaaag	aaaaattatt	1140
tttttggttt	gggggtatta	ggtagtggtt	ttagaagatt	tttttcgaat	tttttaaaga	1200
aaacgttatt	aataatattt	tttttttcgt	ttttaaaata	agtttttggt	ttttcgcgtt	1260
agaataagaa	gtagaaattt	tttttttgga	gttaagaatc	gtgtcgggtt	tatttagcga	1320
ttttaagggt	aggttttttt	cggagttgtc	ggttgagggt	ttttaattta	gttttttagc	1380
gttatttttt	cgatgtttta	tattaaattg	gaatttttag	tgttcgggaag	aaagggtttt	1440
ttggattaat	cgggttttta	tagtagtttag	atttataggt	tagtttttcgg	ggtttttagtt	1500
ttaggggatt	tttatagttt	tatttagagg	tttggggaga	tttttaggatg	atcgatgggg	1560
taacggatta	taattcgtac	ggcgtacgtt	ttttagttac	gtattttttt	tttcgcggta	1620
gggcgattat	aggttttttc	gtttgagcgt	tcgggcgcgt	cgtttattta	gtatcgtaga	1680
gtgtattagt	tttgaatttt	gtttttaata	tttttttagtt	tttttaagtt	cgcgtaaggg	1740
gatggttagga	agcgtttcgg	gattataggt	tcgtcgggtc	gtatatatttg	tattttggaa	1800
ggggcgagga	tgggaagagta	tcgggtgattt	tttttagtat	tgtttttttt	ttgggttaga	1860
gtaaatgaga	taaatttttg	aagggtttta	tataagttat	agggttggtta	ttttggaatg	1920
aggagaaaat	gtagttagcg	ggatttggtt	ttttgttggt	aattaaagagt	tgggaaaacg	1980
aatgggattt	ttatattttt	agatttttaga	ttttaaaagg	gggtaataaa	cgaatcgttg	2040
tttttttatt	tttaatttat	ttagttattac	gcgtataggg	gatgggggttg	gggtaatttg	2100
tttttttttag	ttttgttagt	aatgttgaat	tgtttaaatt	cgagttgggt	gcgggtgttc	2160
gtttggggtt	ttgggtggtg	tatatgtatt	tttaatgcgt	taagataatt	ttttcggagg	2220
ttattttttt	tttttagttga	tttttttaga	ttgggtcggt	tttttttata	taagagtatt	2280
gggtgaagta	tagtatatat	taattttta	gatagagaat	ttgttcgggt	gttttagtgag	2340
atatttttatt	ttttagaatg	attttattat	tttaagatat	tttttagtttg	aagttgtttt	2400
gtttggggtt	ttatagagtt	ggaagttttt	ttttttttta	ttagaagcgg	tttaattagac	2460
gttggttgga	aaaatttttg	tttcgtattg	tttaagagtt	ataaattttt	atgtttgtat	2520
ttggggaggg	ggtttgattt	tcgcgggttc	gagcgtttag	cgggggttatt	ttcgcggttcg	2580
cgcggg						2586

<210> 13
 <211> 2501
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 13

ttgttggtata	gaatattttta	ttatttaggt	attatgttga	gtatttaata	gttttttttt	60
ttgttttttt	tttttttttt	attttggtatt	ttggagttaa	ttatagtgtt	tgttgttttt	120
ttgtttgtgt	tataagtttt	tattatttag	ttttatttta	taagtgagaa	tatttagtat	180

ttggattttt	gtttttgtat	tagttttgtta	aggataatag	tttttagttt	tatttatgtt	240
tttataaaaag	atatgatttta	gttttttttta	atgggttgat	taaatgaagt	tttaaagata	300
taataataaat	attaattttt	tttttattat	aaaaattttt	tgttgaattt	gatttatatt	360
aaatttaata	gttttgtttt	atgaaaagatt	tttttgataa	atttgatagt	tgatggaata	420
ggagaagttg	tttgttatgt	ttaaagttaa	taagagatta	atatttagaa	taaatggaga	480
tttgtaaatt	aatagaaagt	aggtagtaaa	gttaaagaaa	atagtttaag	gtatagttat	540
taaaagggaat	gtgattatgt	tttttgtagg	gatatgggtg	gagttggaag	ttgttagttt	600
tagtaaattt	atataggaat	agaaaattag	tgagattgta	tggttttatt	tataagtggtg	660
agttgaataa	tgagaatata	tggttatatg	tggttgatta	atatatattg	gtgtttgttg	720
agtggggtgt	tggggaggga	gagtattagg	aagaaatagt	aagggatatt	gggtttaata	780
tttgggtgat	gggatgattt	gtatagtaaa	ttattatggt	gtatatattt	atgtaataaa	840
tttgtatatt	ttttatatgt	attttagaat	tttaaataaa	agttggatgg	ttaggtgtgg	900
tggtttatgt	ttgtaatttt	agtatttttg	gaagttgagg	tgtgtagatt	atttaaggtt	960
aggagtttga	gattagtttg	gttaatatgg	tgaatttttg	tttttattaa	aaatataaaa	1020
attagttaga	tgtggtatgt	atttataatt	ttatttattt	gggagggtga	agtagaattg	1080
tttgaatttg	agaggtggag	gttgtagtga	gttgttgaga	ttgtgttatt	gtattttagt	1140
ttgggttata	gtgtgagatt	atgttataaa	ataaaataaa	ataatataaa	ataaaataaa	1200
ataaaataaa	ataaaataaa	ataaaataaa	ataaaataaa	ataaaaaaat	aaaaataaat	1260
aaaataaaaat	aaagtaattt	tttttttttt	aagtggtttt	tatttttttt	ttttgttttg	1320
tgaagtgggt	gtgtaagttt	tgggattgta	tggtttttag	ggaatttttt	tttgtgatgt	1380
tttgggtgtg	tagtttggtg	tgtatatatt	gttgtgggtt	tttttttgtt	gtttgtttat	1440
tttttaggtt	ttgttgggga	tttgggaaag	agggaaaggt	tttttttggg	agttgtgtgg	1500
tgattttggg	gatttttaggg	tgtttttttg	tggttgatgt	ttggggtgta	gtgggtgttg	1560
gggttggggg	tgggtgggag	ttgtgggatt	ttttagaaga	gtggttgggt	ttgtgattta	1620
gtattggggg	ggagtggggg	gggatttatt	ttataagggt	tggagggttg	gagggttttg	1680
ttggagtttt	gttgttgtag	tttttgttat	tagtgagtat	gtgtgggttg	tgtttttggg	1740
gatgggggtt	agagttttta	gtatgggggt	aatttgtagt	attagggttg	ggtttttggg	1800
aggggttttt	gtttattttg	agatttggga	tgggggttta	ggggatttag	gatgttttta	1860
gtgttggttag	tggtttttag	gggggttggg	gtgttttggg	gagggatggg	attttggggg	1920
tggggagggg	gggtagattg	tgtttattgt	gttttgggtt	tttttttggg	gttttagtaa	1980
attttttttt	gtttgttgta	gtgttggttt	atattgtggt	ttatttttta	gtttgaggta	2040
ggagtatgtg	tttggtaggg	aagggaggta	gggggtgggg	ttgtagttta	tagttttttg	2100
tttatttgga	gagatttgaa	tttttttatt	tttttgttgt	gtgggtttta	ttttgggttt	2160
tttttttgtt	ttttgttttt	tttgttatgt	ttgttttttg	tttttagtgt	gtgtgaaatt	2220
tttggaggaa	tttgtttttt	tgtttttttt	ttgtattttt	gatttttttt	tgggttgttg	2280
tgaggtggag	tgggtttggg	ttttatatatt	tgtatttttt	tttttttgta	gggttgtgtg	2340
tgggtttgtg	tatgttggtg	gtagattagg	gttagagttg	gaaggaggag	gtggtgattg	2400
tggagatgtg	gtaggagggt	ttattttaaag	ttttttgtgt	aagtgattat	gtttgggtaa	2460
ggggaggggg	tgttgggttt	taggggggtg	tgattaggat	t		2501

<210> 14

<211> 2501

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 14

gatttttagtt	atagtttttt	aaggtttagt	attttttttt	tttgtttggg	tatgggttatt	60
tatgtaggag	gttttgagtg	agtttttttg	ttatgttttt	atgggtatta	tttttttttt	120
ttagtttttg	ttttgatttg	ttagtagtat	gtgtagggtt	gtgtagtggg	ttgtggggag	180
ggagaagtat	gagatgtggg	gattcgggtt	attttgtttt	gtagtaattt	ggggaggggt	240
taggagtgtg	gggaggggaat	agggaaatag	gtttttttga	agattttata	taatatctgg	300
gtggggagta	ggtatggtgg	gagaggtggg	gaatagggaag	gaggtttggg	gtaaaagtta	360
tatgatggag	ggataagggg	gtttggattt	ttttgggttg	gtgaggggtt	gtgggttgta	420
gttttagttt	ttgttttttt	tttttgttag	atatatgttt	ttattttgaa	ttggggaata	480
gattatgggt	taggggtggt	ttgtagttaa	taaagaaaag	tttgttggag	tttgggggag	540
gatgttaagg	tgtggtgagt	gtagtttgtt	tttttttttt	gtttttgggg	ttttattttt	600
ttttgaggtg	ttttgggttt	tttgaaagtt	gttaattggt	ttggggatgt	tttgggtttt	660
ttaggttttt	gttttgggtt	ttgaggtggg	tgaggagttt	tggttgggag	ttgggtttga	720
tgttgtgggt	tgggttttat	ttgggagttt	tgagttttat	ttttggggat	gtgggttgtg	780
tgtatttatt	ggtggtgaa	attgtggtgg	tgaattttta	gtgaagggtt	tgtgggtttt	840
gagttttata	aggggtggtt	tgtttttgtt	tgttttagtg	ttgagttatg	gtgttgggtg	900
ttttttttga	gggtttttgt	gattttttgt	ggtttttagt	ttgggtggtg	ttgtattttg	960
ggttgttggt	ttagaggggt	gttttggagt	ttttggagtt	gttgtgtagt	tgggtgggga	1020
agtttttttt	tttttttttag	gttttttagt	gggttttagg	agtaaataga	tagtaggaag	1080
aggattgtag	tgaagtgtgt	gtagtgaatt	ggtgtgttgg	gatattgtgg	ggggaaattt	1140
tttaagattg	ttgtgatttt	ggagtttgta	tattttgttt	atagggtagg	ggagaggggt	1200
ggaggttgtt	tagaggaaag	gaaattgttt	tattttattt	tatttttttt	tatttttttt	1260

ttttatttta	ttttatttta	ttttatttta	ttttatttta	ttttatttta	ttttgtgtta	1320
ttttatttta	ttttatgatg	tagttttatg	ttgtgggtta	ggttggagtg	tagtgggtgtg	1380
attttgggtg	ttttattgtaa	ttttgttttt	ttgggttttaa	gtaattttgt	tttagttttt	1440
tgagtaggtg	gaattatagg	tgtgtgttat	atttgggtga	ttttgtatt	tttagtagag	1500
atggggtttt	attatgttgg	ttgggttggg	tttgaatttt	tgattttagg	tgatttgtat	1560
gttttgggtt	tttaaagtgt	tgggattata	gggtgtgagtt	attatgtttg	gttgtttaat	1620
ttttatttga	agttttgggg	tatatgtaga	ggatgtgtag	gtttgttata	taggtgtgtg	1680
tgttatgatg	gtttgttgta	tagattattt	tattatttag	gtattaaagt	tagtattttt	1740
tagttatttt	ttttgggtatt	tttttttttt	agtattttgt	ttaataggta	ttagtgtgtg	1800
ttgattgttg	ttatgtgatt	atgtgttttt	attgttttagt	ttttatttat	aagtgtgatt	1860
atgtgggttt	gttgggtttt	tgtttttgtg	tgagtttgtt	gagggttaagt	gttttttagtt	1920
ttatttatgt	ttttgtaaag	gatatgatta	tgtttttttt	agtggttgtg	ttttagggtta	1980
ttttttttgg	ttttgttgtt	tattttttgt	tgatttgtag	atttttattt	atttttagata	2040
ttgatttttt	gttgggtttta	gatatgatag	atagtttttt	ttatttttatt	aattgtttaag	2100
tttgtttaag	gagtttttta	tgaataaaaa	tttgttaatt	taagtgtaat	taaatttagt	2160
aagggatttt	tgtgggtggg	aagaggttgg	tgtttatgtt	gtatttttaa	aattttattt	2220
aatgtagtta	ttaaaagaa	ttagattatg	ttttttgtgg	gaatatggat	ggagtttagag	2280
gttattattt	ttagtaaat	aatgtaggaa	tagaaattta	aatattggat	gtttttattt	2340
gtaagtggga	gttaaatgat	gagaatttat	aataataata	aggaaataat	agatattgtg	2400
gttgattttta	gggtgttagga	tggggaggaag	gagaggagta	gaaaagagaa	ttattgggta	2460
tttgggtataa	tatttgggtg	atgaaatatt	ttgtataata	a		2501

<210> 15

<211> 2501

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 15

tttgtaaatg	gagatatttt	tattattttt	atagtattat	atgtttttta	agtttgtatt	60
tatatatttg	gtgataaatg	aaggataaga	ttttttttta	tttttgtgag	gatgattata	120
gtatgattgg	atgggtttgt	tatgattttt	attttttttt	gtgtttttat	tattgtttta	180
ttaattttag	ttttttttta	tagggtagta	tagaatttaa	ttagtagaaa	gagatttagt	240
tatgtagatt	agagatttgt	ttaatgtatg	gtatgtaaga	attaggaagg	aaagtttttt	300
gtttaaatat	taataggttt	tttttttaaa	gtaattatta	ttttttaaat	ttaatttata	360
aggtgatagt	atttttaaat	taattaaatt	agaatttttg	gttggataat	tttaaatatg	420
atttattagt	attttttatt	aattattggt	tttttagggt	tttaagttta	tttattagga	480
attttatttt	taattatttt	ttattaaatt	tagttgtaaa	taagagaata	tttaaagggt	540
gaggaatttt	tagtggtaaa	gttttgttta	tgttaagtaa	taaaggataa	gttagttttt	600
gttgtgatta	ttttgttgta	ttgataagtt	atgtattttt	atttaaggat	ttaaattttt	660
atttttttta	agaattgggt	taaaattgat	aaattaaatt	tatttatggt	ttattgatta	720
aagggtgttg	tataataagt	ttttgttatg	tttagtagtt	ggatttatag	tgtagaaat	780
ttataaattg	ttgatttttt	tttttattat	attgtgaaaa	ttgtttttta	aatgtaatta	840
attttaaaaa	tttaaatagta	ttgtgggttag	gtgtgggtgg	ttattattgt	aatattaata	900
ttaggtatag	gtgaggggat	tgaggttagg	atattgaaat	tagtttggga	aatatatgga	960
gatttgggtt	ttggaaaaat	aattagtttt	gtgtgggtgg	gggtgtgagg	ttttgggttaa	1020
ttggggaggt	atagttagtt	atgatgatat	tgtattatag	tttgtgtgat	ggtttatggt	1080
agtaagtttt	ggagtatatt	aaataagttg	tgttgggtat	tttattttatt	ggagagtgat	1140
tagtgattga	tgtttattta	tagtgattag	agatgtatgt	tttgatagta	gtataaattt	1200
agtaggtgtg	aataaatggt	aaagagaaat	tgggtaaaata	agtattatgg	tttttttagtt	1260
gagaaagtgg	gggtttttaaa	aagggttttt	tgttgataga	aagggatggt	taattattga	1320
aattgtagag	gggtgtggtt	tgggtgttga	gtgtgttagat	tatatattatg	gtggtgattg	1380
ttttgtgttt	gggtgtgttt	gtataggtta	tgggtgtttt	gattatgttt	tttaggaata	1440
tttttagtat	ttttgtagtt	ttttttaga	tggaggttga	gatgtgtttt	atgttgttgt	1500
ggtagtaag	gtgttggatg	gttgggttgg	tgatgttttg	gatattgttg	tgtagtattt	1560
tatgggtgtg	tttagtgttg	tttttgttaa	gatttttttt	gtttttgttg	tgggttagata	1620
tgatgagtaa	gaggagtttt	atttaagtgt	ttgtgaggat	tttgggttga	ggtagtggtt	1680
ttatatgata	gttgggtggat	tgaattgaga	atttgaaaga	agttgggtgg	aagttttgtt	1740
ttggtggggg	aggggaaatt	taaagggtta	aattgaaata	gggggaaaaa	aaaagttagt	1800
tttttgtttt	tgtgttttga	attttgtaat	gtgtatagta	ttttgttatt	atgttatgag	1860
gttttaaaaa	attgtttttg	aatgtagaag	atatatatta	atattgtggg	aaatataaga	1920
aaggataaga	aattaagaaa	ttataatggt	attttattat	atagggttagt	taattatgta	1980
ttttgtagag	tagttgtata	tattttttta	agaaaatgta	tatagtgttg	tatatggagt	2040
tttgaatttt	ttttatattg	attataaatt	aattaaattt	tattaaagag	ataaaagtga	2100
tgttttgggt	ttttatgttt	ttaggaattt	ttaatagtta	taattagttt	tttagttaatt	2160
ttttaattgg	ttgtattttt	aaaataatgt	tttttatatt	taataataat	gtattttttt	2220
tttatatttg	ggattaatat	tgaattttat	gattttatta	tattaaaatt	taaattttat	2280
tatattaata	tttaaaattg	tattagaggt	tttatgattt	ggtattatgg	gtttttgtat	2340

tatttttttt	ttaaattttt	taatttggtt	tattaagggt	tttggataat	tttagagatt	2400
ttttgtgaag	tttgaataaa	atttttttga	gatttttgata	attgtattag	ttttaggatt	2460
taattggaat	agaattaaaa	tttttaaaat	aagtttttat	a		2501

<210> 16
 <211> 2501
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 16

tataagagtt	tgttttaagg	attttaattt	tatttttaatt	aagtttttaa	gttaaatgtaa	60
ttattaaaa	tttgaagaga	ttttatttaa	attttataaa	aggtttttta	agttgttttag	120
aaattttggt	gaaatagatt	aggaaatttg	gaaaggaaat	aatgtggaga	tttgtagtat	180
taaattatga	gattttttaat	ataattttta	atattaatgt	aataaaattt	aaattttggt	240
gtaataaaa	tataaatttt	aatattgggt	ttaagtatag	agaaaaagta	tatttatggt	300
gaatgtggaa	aataattattt	ttaaaatata	gttgattaaa	aaattgttgg	ggaattgatt	360
ataattattg	ataattttta	agaaatatag	atattaaaa	attattttta	ttttttta	420
agaaattggt	taaattataa	ttaatatag	gaggttataa	aattttatat	ataatattgt	480
atatattttt	ttggaaaaat	atgtgttaatt	gttttgtaaa	atatatgatt	aattagtttg	540
tgtgatggga	taatatgtga	gtttttta	tttttgtttt	tttttgtatt	ttttatagta	600
ttgatgtata	ttttttgtgt	ttaaaagtaa	ttttttaaag	ttttataatg	tggtataaaa	660
atattatgta	tgttataaaa	tttagaatat	ggaaataaga	agtttgtttt	tttttttttt	720
ttattttggt	ttggtttttt	agattttttt	ttttttattg	gggtgggatt	ttttgttgat	780
tttttttagg	tttttagttt	ggtttgttaa	ttgttgtata	aaggtgttgt	tttaggttag	840
agtttttata	aagtgttggg	tgagattttt	tttgtttgtt	atgtttgggt	gtggtaaagg	900
tggggaaggg	tttggtaaag	gtggtgttaa	gtgttattgt	aaagtattgt	gtgataatat	960
ttagggtatt	tttaagttgg	ttatttgggt	tttttgttgt	tgtggtgggt	tgaagtgtat	1020
ttttgggttt	atttatgagg	agatttgtgg	gggttgaag	gtgtttttgg	agaatgtgat	1080
ttgggatgtt	gtgatttata	tagagtattg	taagtgttaag	atggttattg	ttatggatgt	1140
ggtttatgtg	tttaagtgtt	agggttgtat	tttttatggg	tttgggtggt	gagtgttttt	1200
ttttattaat	aaaagggttt	tttttaggggt	tttttttttt	tagttgagga	gttgtgatgt	1260
ttgtttgttt	agtttttttt	tattattttg	ttgttgttgt	tgagtttgtg	ttgttattgg	1320
agtatgtgtt	tttagttgtt	gtaagtaggt	attagttatt	aattgttttt	tagtaaataa	1380
aatattttaat	ataatttgtt	ttagggtgtt	tagagtttat	tgatatgggt	tggttgtgtag	1440
attgtagtgt	agtgttatta	tgggtttattg	tagttttttg	attagttgga	attttgtggt	1500
tattattatg	taaggttaat	tattttttta	aagattgggt	ttttgtgtgt	tttttagggt	1560
agttttgata	ttttgggttt	aattttttttg	tttatgttta	atgttgggtat	tatagtagtg	1620
agttattatg	tttgggttatg	atattgttga	ggtttttaggg	ttagttatat	ttaaggggta	1680
atttttgtag	tgtagtgggg	aggaaagtta	agtagttata	ggtttttgggt	gttgtgaatt	1740
taattgttga	atatagtaag	aattttattat	gtaataattt	ttaattagtg	gattgtaaat	1800
aagtttagtt	tattgggtttt	ggtttaattt	ttgagaaagg	tgagaatttg	aatttttgag	1860
tagaaatatg	tagtttatta	gtataataaa	gtgattataa	taaagattaa	tttatttttt	1920
gttattta	gtgggtagag	ttttatttgt	gagaattttt	tagtttttga	gtgttttttt	1980
atttataatt	gaagttgata	agatgggtatt	aaaagtgaga	tttttagtaa	gtaaatttaa	2040
aggtttgaag	gatttagtgat	taattgggaag	tgttaataag	ttatatattga	ggttatttaa	2100
tttgagattt	tgatttaatt	ggtttaagg	tattattatt	ttgtgggtta	gatttgaaaa	2160
ataataattg	ttttaaggaa	ggaatttgtt	ggatttttaa	taaaaaattt	ttttttttga	2220
tttttatatg	ttgttattta	gataaatttt	tgggttatat	ggttggattt	ttttttgtta	2280
gttaaatatt	gtgttatatt	gtgaaaaaga	attgagatta	ataaaatggt	agtgagaata	2340
tagggaaaga	taaaaattat	agtaagttaa	tttagttatg	ttgtagttaa	ttttataagg	2400
atagggaagg	attttgtttt	ttatttatta	tttaaaagtgt	gagtataaat	tttaaaaaata	2460
tatgatatta	taggaataat	gaagatgttt	ttatttgtaa	a		2501

<210> 17
 <211> 2251
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 17

ggattttgag	ttgttttttag	gtagtttagga	gtggtttttag	tggtagtgtgt	aaatttttagt	60
agttgtttgt	gttgtttgtg	gttgggggtg	agggtagtta	gagttgggga	ttaagggtttt	120
gtgttatattg	tgtgtatagt	tttatatttg	aatgttgttt	ttttgtagat	gagattgggtg	180
ggtattgtaa	agttgggatt	tgtttttgaa	ggaaaaaaa	tagtgagtaa	gaaatttagt	240

attat	tttt	tttt	tttt	tttt	gaaagt	300
tttt	tttt	tttt	tttt	tttt	gaaagt	360
tttt	tttt	tttt	tttt	tttt	gaaagt	420
tttt	tttt	tttt	tttt	tttt	gaaagt	480
tttt	tttt	tttt	tttt	tttt	gaaagt	540
tttt	tttt	tttt	tttt	tttt	gaaagt	600
tttt	tttt	tttt	tttt	tttt	gaaagt	660
tttt	tttt	tttt	tttt	tttt	gaaagt	720
tttt	tttt	tttt	tttt	tttt	gaaagt	780
tttt	tttt	tttt	tttt	tttt	gaaagt	840
tttt	tttt	tttt	tttt	tttt	gaaagt	900
tttt	tttt	tttt	tttt	tttt	gaaagt	960
tttt	tttt	tttt	tttt	tttt	gaaagt	1020
tttt	tttt	tttt	tttt	tttt	gaaagt	1080
tttt	tttt	tttt	tttt	tttt	gaaagt	1140
tttt	tttt	tttt	tttt	tttt	gaaagt	1200
tttt	tttt	tttt	tttt	tttt	gaaagt	1260
tttt	tttt	tttt	tttt	tttt	gaaagt	1320
tttt	tttt	tttt	tttt	tttt	gaaagt	1380
tttt	tttt	tttt	tttt	tttt	gaaagt	1440
tttt	tttt	tttt	tttt	tttt	gaaagt	1500
tttt	tttt	tttt	tttt	tttt	gaaagt	1560
tttt	tttt	tttt	tttt	tttt	gaaagt	1620
tttt	tttt	tttt	tttt	tttt	gaaagt	1680
tttt	tttt	tttt	tttt	tttt	gaaagt	1740
tttt	tttt	tttt	tttt	tttt	gaaagt	1800
tttt	tttt	tttt	tttt	tttt	gaaagt	1860
tttt	tttt	tttt	tttt	tttt	gaaagt	1920
tttt	tttt	tttt	tttt	tttt	gaaagt	1980
tttt	tttt	tttt	tttt	tttt	gaaagt	2040
tttt	tttt	tttt	tttt	tttt	gaaagt	2100
tttt	tttt	tttt	tttt	tttt	gaaagt	2160
tttt	tttt	tttt	tttt	tttt	gaaagt	2220
tttt	tttt	tttt	tttt	tttt	gaaagt	2251

<210> 18
 <211> 2251
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 18

tgtag	tttat	agttt	atagt	atagg	aggat	60
ttatg	tttaa	tagagg	gggtt	tttga	ttagg	120
agggg	taagg	gttgg	ttttg	aattt	ttggg	180
gatgt	agggg	taaa	agttt	gttag	ggatt	240
ttgtt	tttga	gaaat	agttg	gggaa	tttat	300
taagg	tttaa	agggg	agttg	gggag	ttgaa	360
aggtt	gagta	gggtt	tttag	tattt	gggat	420
tattt	gggag	gtgga	taata	tattt	ttggg	480
gaagt	ttttg	tggtt	ttgaa	agttt	gtggt	540
gtttt	gggag	gtaatt	gagg	tggtt	ttagg	600
gtggt	gttat	aattt	gggt	gtggt	ttatg	660
gatgt	gggtt	agaa	aggg	gtttt	atgag	720
tggtt	gttat	ttggg	gttat	gtatt	ttggg	780
gttat	gagaa	tttat	gttat	aaagt	ttggg	840
gagtt	gtgag	tttat	gttat	tagaa	tatgg	900
ggtag	gttat	tttat	gttat	aggtt	ggtaa	960
gagta	aaagt	atatt	gttat	ttggg	ggttt	1020
gtatg	gggtt	atggg	tttat	taaag	aatag	1080
tttat	atatt	aggtt	tttat	ttttt	ttgtt	1140
tatgat	gggtt	tttat	tttat	gaagt	attat	1200
tggtt	gttat	aggtt	gggtt	gttat	ttgat	1260
gatag	gggtt	gttat	gggtt	gttat	ggtaa	1320
gagata	aaagt	gttat	gggtt	gttat	gttat	1380
aatgat	atatt	aggtt	tttat	tttat	tttat	1440
tattat	gttat	atatt	tttat	tttat	tttat	1500
tggtt	gttat	atatt	tttat	tttat	tttat	1560

tattttattg	ttaagatgaa	taagatttat	tttaagggaa	aagtagtttg	ttagtgttat	1620
ttttgataga	ttttgaatta	ataaaaatggt	tattgaaaat	aagttataaa	agtaatttgg	1680
agagtgtgtt	tttttttttt	tgggttttttt	tgggtgaggt	tgtattttttg	tgttttttat	1740
ttgttgattt	taatgttagt	ttaatgttat	tttatttgta	gtggtagtag	tggtggtagt	1800
tttatgagtt	tttagggtta	gttttttagat	gtttttggatt	tgtttttaat	ttgtttttat	1860
tttgttttgt	ttagtttgga	aattgttaaa	ttttttggttt	tttaaagttg	gttttagtggt	1920
ttgttgattt	ttggatattg	agggttagagt	tgtagtttt	taaaaatttg	ggaaataaga	1980
ggtgtagtgg	gatgggttag	tgaagaatgg	tggtggattt	tttatttggt	attttttttt	2040
ttttaagat	gagttttagt	ttttagtagt	ttgttggttt	tgtttggtggg	aggatagtg	2100
ttaggtgtga	ggttggtgt	gtaggtggtg	tggagttttg	gttttttagtt	ttggttggtt	2160
tttgtttttag	ttatgggttag	tatgggtggt	tattggagtt	tgtggttggt	gttgaggttg	2220
tttttggtta	tttgggggtg	gtttgggggtt	t			2251

<210> 19
 <211> 2586
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 19

tttggtggtg	tgtgggagta	gttttggttg	gtgtttgtag	ttgtgggagt	taagtttttt	60
ttttaggtgt	aggtataaaa	gtttatgggt	tttgaataat	gtggggtaga	ggttttttta	120
agtaatgttt	aattggttgt	ttttaattaa	ggaaagagag	gttttttagt	ttatggtaat	180
ttaagtaggg	tagtttttag	ttaaagggtat	tttagaataa	taagattatt	ttaagaaatg	240
gaatgtttta	ttggatattt	gaatagggtt	tttggtattg	gaattggtgt	gtattgtatt	300
tttaattagta	tttttgtgtg	gagggaggtg	atttagttta	ggaaagttaa	ttatagaaag	360
aggtgatttt	tgaaggatt	gttttagtgt	tattagaata	tatgtgatta	tattaaaagt	420
ttaggtggat	attttagtgt	agtttggatt	tggataattt	aatattgttg	gtagaattga	480
agggataaag	ttattttaat	tttatttttt	gtatgtgtag	tggtgagtga	gttgggggtg	540
ggaggatagt	ggtttgttta	ttgttttttt	ttaaaatttg	agatttgaaa	atatggagggt	600
tttatttgtt	tttttagttt	ttgattgtta	ataaaaaaat	aaattttgtt	ggttatatatt	660
tttttttatt	ttaaaatagt	aatttttatgg	tttgatttaa	gttttttaga	agttttattt	720
atttgttttg	ggattaggag	ggaataatgt	taggaaaagt	tattggttgt	tttttatatt	780
tggttttttt	aggggtgtag	atgtgtgggt	tgggtgggtt	gtgatttttg	aatgtttttt	840
gttatttttt	tgtgtgaatt	tgaagggtt	gggaggtgtt	gagagtagag	tttaggggtg	900
gtgtattttg	tgggtgttag	tgggtgtgtg	gtttgggtgt	ttaggttggg	ggattttag	960
ttgttttatt	gtggagggga	aaatatgtag	ttggaggggt	tgtgttgtgt	gggttgtgat	1020
ttgttatttt	attggttatt	ttgggttttt	tttaagtttt	taggtagggt	tgtgagagtt	1080
tttttagagtt	gaagtttttg	aggttgattt	gtgggttttg	ttgttatggg	aatttggttg	1140
gtttaaagaa	gttttttttt	tgggtatttt	gaatttttagt	ttagtgtggg	gtattggggga	1200
agtggtgttg	gggggttggg	ttgggggatt	ttagtgggt	gttttgagga	gggtttattt	1260
ttgggggtgt	tgggtgaggt	tgggtatgatt	tttggtttta	aaaggaaaagt	ttttgttttt	1320
tgtttttggg	ttagagagtt	tttatttatt	ttgagagttg	agagagaaat	gttattgggt	1380
atgttttttt	tggaaagttt	gagaggggtt	ttttggatat	attatttagt	gttttttaaat	1440
tagagaagta	gttttttttt	ggtgtttggg	tttagaagtt	gttatttatt	tagtttatgg	1500
tttgaaatta	gtatgggaag	tgttggggta	aggtttttgt	ggagattaga	ggtttgtttg	1560
ttggggaggag	tttttggggg	atggggattt	tatttttttt	tttgttttgg	tttttatttg	1620
ggatgttttt	ttaggagttt	agaaaagatga	tttattatat	ggttttggga	tagagtagtg	1680
tgtttaattt	ttaggggaatt	ttgtgtgttt	ttttgaggtt	ttagtttttt	aaggtattgt	1740
tgtttgtttt	ttttttttta	gattgaaatt	ggggaagagt	gtgggtgttt	ttttgttttg	1800
atgagtttgt	tttttttaaat	gtttattttg	gttgatttag	agtatttggg	aaattttgaa	1860
aggtgttttag	gttttatata	gtagtgtttt	tttatttagt	ttttgttttt	gggttttttt	1920
aagagagttt	ttattttatg	ttttgggttt	tttttgatgt	tgggtttttg	aggtagggtat	1980
ggagtttttt	tgaagtagt	tgtttatttg	tgtttttttg	gtgtaaagtt	agagtttatt	2040
ttgttggggg	aaggggaggt	agaaaagatt	atagttggga	aagtgtgttt	tttgttttgt	2100
ttttaaaata	tgttttaaga	ttgttattgt	gattgttagg	agagttatta	atgttttaggg	2160
gttataaagg	aaatttttgaa	tttttgggtt	tttttaaat	tttaggtttt	taaaaattta	2220
gtggggggtt	tttgggggtt	ggatttaggt	tgttatttgt	gggaggattt	tgtttagtat	2280
ttttttatta	atattttatg	aaggtaggtt	tttgtttttt	ttggagtttt	tttttttgga	2340
atgtttttta	attttgggtt	atttattttt	ttgtgagtta	ttttagggtt	ttgtgggttg	2400
ggaagagatg	tgtttaatttt	gtgggtttgt	gtgtagtttt	tagttgtaaa	gtgttgttaag	2460
tgattttttt	gatgggtttt	tttgattgaa	gagtttggga	attaaagaga	aaaaaaataa	2520
ttttattttt	aaaagaataa	gttattattg	tgggtgatatt	gtggtggagg	attttgggtga	2580
tgggggt						2586

<210> 20
 <211> 2586
 <212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 20

attttattgt	taaagttttt	tgttatagta	ttgtttagt	gatgatttgt	ttttttgaaa	60
ataaagttat	tttttttttt	tttgggtttt	gagttttttg	gttggaaagg	gttgttaggg	120
gggttatttg	tagtattttg	tggttaagga	ttgtgtgtag	atttgtggag	ttgatgtgtt	180
tttttttggg	ttatagagtt	ttaggatggt	ttatagggga	gtgagttagt	tagagtttgg	240
gaatattttg	agaaaagagg	tttttagaga	ggtaggagtt	tgtttttgtg	aaatattaat	300
aaagggatgt	taggtgaggt	tttttttagtg	gtgttagttt	gaattttagt	tttaagagat	360
ttttattagg	gttttaggaa	tttggggggt	tgggaagggt	tgagggttta	gaaatttttt	420
tatagttttt	agatgttgat	agttttttta	ataattgtga	tgatagtttt	gaggatatgt	480
ttaggaataa	ggtgaaaagt	gtattttttt	aattgtgatt	ttttttattt	tttttttttt	540
taataaagta	aattttaatt	ttatattaaa	ggggtataga	taggtaattg	tttttagagg	600
gatttttgtg	ttattttggg	gatttgatat	tgaagaaaga	ttgaggtagt	aggtagagat	660
tttttttgaaa	aaattttaag	atagaggttg	agtagggaga	tggtgttgtg	tgaggtttgg	720
gtattttttta	gagttttttt	gatgttttgg	tgtagttaga	gtagggtgtt	ggggagggtga	780
atttatttggg	gtaaagaagt	gtttatat	tttttttagtt	ttgggttagg	gaaagaagaa	840
tggatggttg	tgttttggaa	agttaggggt	ttagagaggt	gtataaagtt	tttttaaagt	900
tgggtgtgtt	gttttgtttt	gggattatgt	agtggattgt	ttttttgggt	ttttatggag	960
gtgttttagg	tgggaattag	agtaagtagg	agaatggggt	ttttattttt	taggggtttt	1020
ttttgatagg	taggtttttt	gtttttgatg	aagtttttgt	ttgggtgttt	ttatgttgat	1080
tttgaattat	gggttgagtg	agtgggtgatt	tttgagttta	ggtatttaaag	aaaaattatt	1140
tttttggttt	gggggtatta	ggtagtgtgt	ttagaagatt	ttttttgaat	tttttaaaga	1200
aaatgttatt	aataatattt	tttttttgt	ttttaaaata	agtttttgggt	tttttgtgtt	1260
agaataagaa	gtagaaattt	tttttttggg	gttaagaatt	gtgttgggtt	tatttagtga	1320
ttttaagggt	aggttttttt	tggagtgtgt	gggtgagggt	ttttaattta	gttttttagt	1380
gttatttttt	tgtatgtttt	tattaaattg	gaatttttagg	tgtttggaa	aaagggtttt	1440
ttggattaat	tgggtttttt	tagtagttag	atttataggt	tagtttttgg	ggttttagtt	1500
ttaggggatt	tttatagttt	tatttagagg	tttggggaga	ttttaggatg	attgatgggg	1560
taatggatta	taatttgtat	ggtgtatgtt	tttttagttat	gtattttttt	ttttgtggta	1620
gggtgattat	aggttttttt	gtttgagttg	ttgggtgtgt	tgtttattta	gtattgtaga	1680
gtgtattagt	tttgaatttt	gtttttaata	tttttttagtt	tttttaagtt	tgtgtgaagg	1740
gatggttaga	agtgttttgg	gattataggt	ttgttgggtt	gtatattttt	tattttggaa	1800
gggggtgagga	tgggaagagta	ttggtgattt	tttttagtat	tgtttttttt	ttgggttaga	1860
gtaaatgaga	taaatttttt	aagggtttta	tataagttat	aggggttgta	ttttggaatg	1920
aggagaaaaa	gtagtttagtg	ggatttggtt	ttttgttgggt	aattaagagt	tgggaaaatg	1980
aatgggagtt	ttatatattt	agatttttaga	ttttaaaagg	gggtaataaa	tgaattgttg	2040
tttttttatt	tttaatttat	ttagtattat	gtgtataggg	gatgggggtg	gggtaatttg	2100
tttttttttag	ttttgttagt	aatgttgaat	tgtttaaat	tgagttgggt	gtgggtgttt	2160
gtttgggttt	ttgggtgtgt	tatatgtatt	ttaatagtgt	taagataatt	tttttggagg	2220
ttattttttt	ttgtagttag	tttttttaga	ttgggtgtgt	tttttttata	taagagtatt	2280
gggtgaagta	tagtatatat	taattttaat	gatagagaat	ttgtttgggt	gttttagtgag	2340
atatttttatt	ttttagaatg	attttattat	tttaaggtat	ttttagtgtg	aagttgtttt	2400
gtttgggttg	ttatagagtt	ggaagttttt	ttttttttaa	ttagaagtgg	tttaattagat	2460
gttgttttgg	aaaatttttt	ttttgtattg	tttaagagtt	ataaattttt	atgtttgtat	2520
ttggggaggg	ggtttgattt	ttgtggttgt	gagtgttttag	tggggttatt	tttgtgtttg	2580
tgtggg						2586

<210> 21

<211> 2308

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 21

ttaggttagga	ggtagttagt	tgggtgatagc	gcgcgtgatt	atagattttt	gattttttata	60
gttttagtag	ttgtttgtag	gtttttttacg	ggatcgtagt	ttgggaggggt	aggagggtttt	120
gtatttgcgg	tgggtttgat	ttgttttttta	tttttttgtt	tgttttttttc	gtgttgggag	180
tcgtcgcgat	acgttttcgcg	gatatttagg	agtttttttta	gttttttgcga	gcgcggatttc	240
gggagagagt	ttttatttaa	cgttgaggtt	attggtttat	agatttttttc	gggtttatttt	300
gcgaagtgtt	tgggtttttt	tatttttagt	tttgttagtag	taataaacgg	gaaggaagtg	360
ggtgacgagg	ataggtagag	atttttttta	cggggtcggt	cgtattttat	ttgtcgtcgc	420
ggggcgggag	ttgcggtcgg	tgtttagttg	tagattcgta	ggcgatcgta	tttttggcgg	480
tttttagttt	aagtaggtag	cggtcgattc	gttttgttcg	agggttttagg	gttgatttag	540

gtgagtttcg	gcggggaggg	cgggcgtttc	ggttttttcg	tttcggaatt	ttgtatcggt	600
tagagtcggt	acgttttttcg	gttgtaggtt	tcgtatttgt	aaaatgggtc	gatgaatttt	660
aagggtcggt	ttagattttac	ggattgtagt	tgtagttggg	ttagaagggg	tggttagggg	720
agtcgatttt	tttttttttt	ttttttttacg	tttaattagga	gttttttattt	gtttgggatt	780
ttttgggggt	atztatgggt	ttatagtcgt	agaattatag	tattgggtgg	ggtttttagtt	840
ttgggggttaa	gtttttgtga	ggtttggtat	tatttttttt	tgttgttttg	tttttttggt	900
tgtttggttg	tttggttttt	ttttttttga	gatagagttt	cgtcttttta	tttaggtcgg	960
attgtagtgg	cgcgattttcg	gtttattgta	agttttgttt	ttcgggtttt	cgttattttt	1020
ttgttttagt	atttttcgagt	agttgggatt	acgggcgtcg	ttatcgcggt	cggttaattt	1080
tttgattttt	ttagtagaga	cgggggtttta	tcgtgttagt	taggatgggt	tcgatttttt	1140
gatttcgtga	tttattttatt	ttgggtttttt	aaagtgttgg	gattataggg	gtgagttatc	1200
gcgttcgggt	tgttttttgtt	ttttttttttt	ttgagatagt	tttatttttg	ttgttttaggt	1260
tgtagtgtag	tggcgtaatt	ttgggtttacg	taattttcgt	ttttcgggta	taagcgattt	1320
ttttgtttta	gttttttttag	tagttgtgat	tataggtacg	tattattacg	ttcggttaat	1380
tttgattttt	tagtagagac	gggggtttttt	tatgttgggt	aggttgggtt	ttaatgttcg	1440
attttaggtg	atttgtttat	tttagttttt	taaagttttg	cgattatagg	cgtgagttat	1500
tatgtttggg	tagttttttg	tttttttaaa	aatttattat	ttagtcgggt	tggtgtttta	1560
tatttgtaat	tttggtattt	tgggaggtcg	aggtgagcgg	attatttgag	gttaggtggt	1620
ttagattagt	ttgggtaata	tggtaaaatt	ttgtttttat	taaaaagata	aaaattagtt	1680
ggaggtggtt	ttttgaattt	aggaggtaga	ggttgtagtg	aattgagatt	atgttattgt	1740
attttaattt	aggaggagga	ggttggtattg	agttaataat	atgttattat	attttagttt	1800
gggcgataga	gtaagatttt	gttttttttta	aaaaaaaaat	agaaaaaaaa	aaaaagaaag	1860
aaatttgaag	taattatgat	taaatgttaa	tatttggttt	tttacgtggg	gggtataggc	1920
gtgttttgga	agtttttatg	ttttttgtat	tttgagatta	tataaaatta	tgtttattat	1980
tttattgtgt	tatatgattt	ttgtagaaaa	gttggtaaat	tttggaatgt	acgaagaaaa	2040
agattataat	agttagttat	atttttgttt	agagagatag	tattttattaa	tttttttttg	2100
tttatgtttg	ttttcggaaa	aatgggatat	tattggtatt	attttataat	tttttaattt	2160
gtattaatat	tttttatatt	ttttaaattg	aaaaagttgg	ttaagtatgg	tggttttaggt	2220
ttgtaaattt	agttaatgtg	ggaggatcgt	ttgagttcgg	gagtttagga	atagtttggg	2280
tatcgtggcg	agatttatatt	gttataaa				2308

<210> 22

<211> 2308

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 22

ttttagtagta	tgtaggtttcg	ttacgatggt	taggttggtt	ttgaattttc	gggtttaagc	60
gatttttttta	tattgggttgg	gtttataggt	ttgagttatt	atgttttggt	agtttttttta	120
atttaaaaaaa	tatagaaaagt	gttaatatata	attaaaaagt	tataaaatag	tgtaaatagt	180
atttttatttt	ttcgaagata	aatataggga	gagaaaaatt	aatggatatt	gttttttttgg	240
gtagggggtat	tattggttat	tataattttt	tttttcgtgt	attttagaat	ttattaattt	300
ttttataaag	attatgtagt	ataataagat	aataaatata	attttggtg	attttaaaat	360
ataggaagta	tagaagtttt	taaaatacgt	ttgtatttat	tacgtgaaga	aatagatggt	420
aatattttggt	tataattggt	ttagattttt	tttttttttt	ttttttttgt	tttttttttg	480
ggggggatag	agtttttggt	tgtaggtttg	gttggaatgt	agtggtatga	tattgggtta	540
gtgtaatttt	tttttttttg	gttgaggtg	agtggtatga	tttttagttta	ttgtaatttt	600
tggttttttgg	gtttaagaga	gtatttttag	tttaattttg	tttttttagt	agagataagg	660
ttttgttatg	ttgttttagt	tggtttggaa	tatttgattt	taagtattc	gtttatttcg	720
gttttttaaa	gtgttaggat	tataggtgtg	agatattata	tcggttaaat	aataagtttt	780
taaaaaaaata	ggagattggt	taggtatggt	gatttacgtt	tgtaatcgta	gaattttggg	840
agggttaggt	gggtagatta	tttgagatcg	ggtagtgag	attagtttgg	ttaatatgga	900
gaaatttcgt	ttttattaaa	aataataaaa	tagtcgggcg	tggtggtgcg	tggttgtaat	960
tatagttatt	agggaggttg	aggtaggaga	atcgtttgta	ttcgggagcg	ggaggttgcg	1020
tgagttaaga	ttgcgttatt	gtatttttag	ttgggtaatt	agaataaaaat	tggttttaaaa	1080
aaaaaaaaaaa	taaaaatagg	tcgggcgcgg	tggtttacgt	ttgtaatttt	agtattttgg	1140
aaggttaagg	tgggtggatt	acgaggttag	gagatcgaaa	ttattttggt	taatacgggtg	1200
aaatttcggt	tttattaaaa	aatatcggtg	attagtcggg	cgcggtggcg	gcgttcgtag	1260
tttttagttat	tcgggggtgt	tgaggtagga	gaatggcgag	aattcgggag	gtagagtttg	1320
tagtgagtcg	agatcgcggt	attgtagttc	gggttggttg	aaagagcgag	attttggttt	1380
aaaaaaaaaaa	aaattaaata	aataaataaa	taaaaaaaata	aaataaatagg	agaagatgat	1440
gatagatttt	ataagagttt	ggttttaggg	ttgaaatttt	attttagtgt	gtgggttttgc	1500
gggttggtggg	ttatgggtgt	ttttaagggt	tttaggttag	atggaggttt	ttgggttggcg	1560
tggagggaagg	aaaaggatga	aatcgggttt	tttggttatt	tttttttagt	tagttgtagt	1620
tgtagttcgt	gggttttagt	cgatttttgg	ggtttatcgg	tttattttat	agatcgggag	1680
tttatagtcg	ggagggcggt	cggtttttgg	cgggtgtagg	tttcggggcg	gggaggtcga	1740
ggcggttcgt	tttttcgctg	gggtttattt	gggttagttt	tgagttttcg	agtagggcgg	1800

atcggtcggt	gtttgtttta	agttgggggc	gttagggagt	cgatcgtttg	cggattttgta	1860
gttgggtatc	ggtcgtaacg	ttcgttttcg	gacggtaggt	gagatgcggg	cggtttcggtg	1920
ggagggggtt	ttgtttgttt	tcgttattta	tttttttttc	gtttgttggt	gttgtaggggt	1980
tgggaatggg	ggggattaa	tatttcgtag	gggtattcgg	ggagatttgt	gagttagtgg	2040
tttttagcgt	gaatgagagt	tttttttcgg	gttcgcggtc	gtagggattg	aagggatttt	2100
tgaatgttcg	cgggacgtgt	cgcggcggcg	tttagtacgg	aggaggtagg	tagggaggtg	2160
aggagtaggt	tagatcggtc	gtaggtgtag	ggttttttgt	tttttttaggt	tgcggttttcg	2220
tggaggggtt	gtagatagtt	gttgagatta	tgagggttag	aggtttgtgg	ttacgcgcgt	2280
tattattagt	tagttgtttt	ttgtttgg				2308

<210> 23
 <211> 2553
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 23

ttggttattt	ttgttttttt	ttagattttt	agagttttta	gtgtagttat	agaggatggt	60
gttggtttta	gttttaagaa	gacgtcgttt	tttttagagg	gttaagtaag	tgggaatttt	120
tttttttatt	tgttttgggt	tttaggtagg	gtttttgggtg	taaggtttgg	ggttggaagt	180
cgatttattt	aggtttaggt	tttggggtag	aattgaaatt	ttttggttat	tgctcggtgt	240
agttttgggag	taggttattg	ttaaaagtgt	gggttttttt	aggatagttt	ttttatgagg	300
tcggtttttt	atttgttggt	tttttataatt	tggtgggttag	ggatgtgggt	ttgggtagaa	360
cgatgatttt	ttatttttgt	tattatggaa	gttatcgttg	tttttttaggt	tagttagtta	420
tttgggttgt	agagtatttt	ttttatgttt	ttcgggtgtt	tttttttttt	tttgttttag	480
tttggttttg	ttttattttg	tttttaggga	gggttatatt	ggagtgggggt	tagggtaggg	540
tttttttttcg	agggagtttt	tttttggttg	tttttaggggt	agttttgtat	agtttttagta	600
tttggcggtat	ttttttttgat	atttttttta	gggatagtta	ggtattttgt	gtgggggtatt	660
taagagagtt	aggttcgtta	gttttttagtt	tttgtagaa	tgtaggtttg	aggggtgagg	720
ggcggggtag	gggtagggat	aggaatttcg	gcgtgttttt	tattcgtaaa	ggtttattga	780
ggtttcagat	tttagttatt	gagttattaa	gttagtttgg	gttaggtttg	ggtgttttgt	840
ttgtaatgga	ggtagagacg	gggtttcggg	gtagttttga	ggatgttggg	tgtatagcgg	900
gggtttcgtc	ggtaggaatt	atttatgttt	ttttttgggt	taagttttgt	ggatgttttag	960
tttgggggtc	cggggagttg	gtaggttagt	ggtagatatt	ggtgggtaga	tttagtgttt	1020
ggtagaatag	gtattaagga	agtgggtgat	ggagggaagt	taagtgtatt	ttaaattttcg	1080
ggtgagttat	tatcgtcggg	ttttttatag	ttgttgaaag	tgagtaatag	tgatgaagggt	1140
ttgtgagttt	ttgcgtgagc	gagtgaatgg	attagtagta	gttttttaggt	tgtggaagag	1200
cgtttttttt	tcgggatggg	gatatttgggt	tatagtaatt	tttaattttt	tatttattta	1260
tcgtttattg	tagaggtagt	cgggggtttt	gttttttgta	ggtaggagtg	aggggtattt	1320
ttgtgatgtg	gtatttttgt	gatcagaggt	atgtgtgatc	ggtgtaaggg	taggaagcga	1380
gttattgggt	tgtattaggc	gtgggggttt	ttgcgagggt	aggatttaaa	gtcgggttgg	1440
tttttcgggt	gtagtatttt	tttttttttc	gaattaggtt	agagttttgg	gacgggaggt	1500
gttttgtaga	ttattttttt	tattaatttt	cgtttttcgt	tttattttcg	cggtagttcg	1560
gtgaattgtc	ggttttttgt	tgtgtatcga	gtggggtagt	gattttgacg	tggcggtttt	1620
tgctcgtttt	gttatcgtta	ttattttcgg	tggttttagtt	ttcgtatttt	ttatttttat	1680
ggaggaatgt	attaggtttt	tttttttgga	tgtatttttt	atttatatgt	ttttaaat	1740
tggtattttt	tggttttttt	ttatttttat	tttttttttt	agggttttag	ataaagggga	1800
agtggttgga	tttttttaaa	gggataggtg	tttttttagtt	tattgttgaa	tttttttttt	1860
taatttttagt	tttttagtta	tagttaatta	gtattagtag	atagtttatg	agtgatattt	1920
atgtaggttt	taggttgtgg	agagtttttt	gggtaggaaa	tagtttttaa	ggttttttat	1980
tttatttagg	tttttagttt	ttttatttgt	ttttttttta	gatttgtggt	ttttggagtt	2040
tggttttttt	gttttttgtg	gacgatata	tagtatttaa	atagtggtag	agcgggacgg	2100
atttttttagt	ttgttttttg	tgtgggtttg	tatttttgatt	tagatatggt	tttttatagt	2160
aggatttagg	gggggtatatg	tgtgtttgcg	ggtttatttg	ggtattcgta	tttgggttat	2220
tttatttttt	agagagaggg	ttttgttgtg	ttatttagtt	ggagtgtagt	ggtgtaatta	2280
tagtatattg	tagtttttaa	tttttgggtt	taagcgattt	ttttttttta	gttttttttag	2340
tagttggggag	tataggattt	attgtatttt	ggtaattttt	ttaataattt	tttaagagat	2400
gggggtttat	tgtgttgttt	aggttgggtt	taaatttttg	gttttaagtg	atttttttat	2460
tttcgttttt	tgaagtgttg	agattatagg	tatgagttat	tatgtttatt	ttagattgat	2520
atttttatat	ttgtttattt	tgggtgggta	ggg			2553

<210> 24
 <211> 2553
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 24

ttttgttttag	ttaggatgaa	taaatataga	aatgttagtt	tgggatgggt	atgggtgggtt	60
atgtttgttaa	ttttagtatt	ttaggagggc	aaggtgggag	gattatttga	ggtagagggt	120
ttgaggttag	tttgggtaat	atagtaagat	tttatttttt	aaaaaattat	taaaaaatta	180
gttaggatata	agtgggtttt	gtatttttag	ttattagggg	ggttggggag	ggaggatcgt	240
ttgagtttag	gagttgaagg	ttgtagtgtg	ttatgattat	attattgtat	tttagttggg	300
tgatatagta	agattttttt	tttaaaaaat	aaaaataaatt	aaatgcgggt	gttttagtga	360
attcgtaggt	atatatgtgt	ttttttgggt	tttgttgttg	gggggtatgt	ttgggttagg	420
gtatagattt	atatagagaa	taggttaggg	ggttcgtttc	gttttgttat	tgtttgggtg	480
ttatgtgtcg	gttatatagg	gatagaagaa	ttaggtttta	aagggttata	atttaggaga	540
gaggtaggta	ggaaagattg	ggatttggat	gagatgaggg	attttaaggg	ttgtttttta	600
tttaggaaat	tttttatagt	ttggggtttg	tatgggtatt	atttatgggt	tgtttgttaa	660
tgtaaattaa	ttgttaattag	ggaattgggg	ttgaggaggg	gagtttagta	gtaagttggg	720
gggatattgt	ttttttaaga	ggatttagtt	attttttttt	tgtttgggag	tttaggggaa	780
ggggtgggag	taaaggggga	gtagaaaatg	ttagggtttg	gaagtatgtg	ggtgaggggt	840
gtatttagga	agggaggttt	ggtgtatttt	tttatggggg	tggggaatgc	ggaggttggg	900
ttatcggagg	tgggtggcgg	ggtagggggc	gtaggagacg	ttacgttagg	gttattgttt	960
tattcgggtg	atagtagggg	gtcggtagtt	tatcggatta	tcgcgggggt	ggggcggagg	1020
acggaagtgt	gtgagggggg	tggtttatag	gggtattttt	gttttagggg	tttaatttaa	1080
ttcgaaaggg	aaaggagtgt	tgtagtcggg	aggtagggte	gattttgggt	ttgttttcg	1140
tagaagtttt	tacgtttggt	gtagattaat	gattcgtttt	ttgtttttat	atcgattata	1200
tatgatttcg	gttatagggg	tggtatatta	taggagtgtt	ttttattttt	gtttgtagga	1260
agttagggttt	tcgtatatatt	ttgtagtggg	cggtaggggt	gtgggggatt	aggaattgtt	1320
gtaattaaagt	gtttttattt	cggggagggg	acgttttttt	ataatttgga	aattgttatt	1380
ggtttattta	ttcgtttacg	tagaaattta	ttaaattttt	ttattgttgt	ttatttttag	1440
tagttgtgaa	agattcggcg	gtgatgattt	attcagaggg	ttgagtgtat	ttggtttttt	1500
ttcgggttatt	atttttttga	tgtttgtttt	attagatatt	aggtttgttt	attagtgttt	1560
gttattgatt	tgtagttttt	tcgcgggttt	aggttgggta	tttataaagt	ttggttttag	1620
agagagtata	agtatttttt	gtcggcgagg	ttttcgttgt	gtatttagta	tttttagaat	1680
tgtttcgaga	tttcgttttt	gtttttattg	tagatagggg	atttaggttt	ggtttaggtt	1740
gatttggttg	tttagtggtt	gggtttcggg	gttttagtga	atttttgcgg	atggagagta	1800
cgtcggaggt	tttggtttttg	tttttgtttc	gttttttatt	tttttaggtt	gtattttggt	1860
aggagttaga	ggttgacggg	tttggttttt	ttgagtgttt	tatatagagt	gtttgattgt	1920
ttttaagaag	gatgttaagg	gaggtgcgtt	aggatttgag	gttgtgtaga	gttggttttg	1980
ggatagttag	agaggaattt	tttcggggga	gagttatgtt	ttgggtttat	tttaggggat	2040
ttttttttga	gagtagggta	ggataaagtt	aggttggggg	aggagaaggg	ggaggtattc	2100
ggaggggtatg	aaaggggtgt	ttttagtatt	agggtggttg	ttgggttggg	agatagcggg	2160
gggtttttata	atgataggag	tggagaatta	tcgtttttat	tagggttata	tttttggtta	2220
ttagggtgtga	agaaatagta	gggtggaggat	cggtttttatg	gggagattgt	tttggaagga	2280
tttatagttt	tggtagtggg	ttgttttttag	gtttagtctg	atagtaatta	aggagtttta	2340
gttttgtttt	agagtttgga	tttaggtggg	tcggttttta	gttttaggtt	ttatattagg	2400
gggttttgttt	ggagttttag	ataagtaggg	aggggggatt	ttattttatt	agtttttttg	2460
aggaagcggc	gttttttttg	ggttgaagtt	aatagtattt	tttgtggtta	tattgggggt	2520
tttgagaggt	taagggaggg	taggggtggg	tag			2553

<210> 25

<211> 2308

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 25

ttaggttagga	ggtagttagt	tggtgatagt	gtgtgtgatt	atagattttt	gattttttata	60
gttttagtag	ttgtttgtag	gttttttatg	ggattttagt	ttgggagggg	aggaggtttt	120
gtattttgtg	ttgggttgat	ttgtttttta	tttttttgtt	tggttttttt	gtgttgggtg	180
ttgttgtgat	atgttttgtg	gatatttagg	agttttttta	gtttttgtga	gtgtggattt	240
gggagagagt	ttttatttaa	tggtgaggtt	attggtttat	agattttttt	gggttatttt	300
gtgaagtgtt	tggttttttt	tatttttagt	ttttagtagt	taataaatgg	gaaggaagtg	360
ggtgatgagg	ataggttagag	atttttttta	tggggttggt	tgtattttat	ttgttgttgt	420
ggggtgggtg	ttgtgggttg	tgtttagttg	tagatttgta	ggtgattgta	tttttgggtg	480
tttagttttt	aagtaggttag	tggttgattt	gttttgtttg	agggttttag	gttgatttag	540
gtgagttttg	gtggggaggg	tggttgattt	gggttttttg	ttttggaatt	ttgtatttgt	600
tagagtttgt	atgttttttg	gtttagaggt	ttgtatttgt	aaaatgggtt	gatgaatttt	660
aagggttggg	ttagattttat	ggattttagt	tgtagtggg	ttagaagggg	tggttagggg	720
agttgatttt	attttttttt	tttttttatg	tttaattagga	gtttttattt	gtttgggatt	780
ttttgggggt	atttatgggt	ttatagttgt	agaattatag	tattgggttg	ggtttttagt	840

ttgggggttaa	gtttttgtga	ggtttgttat	tatttttttt	tgttgttttg	tttttttgtt	900
tgtttgtttg	tttgggtttt	tttttttga	gatagagttt	tgttttttta	tttaggttgg	960
attgtagtgg	tgtgattttg	gtttattgta	agttttgttt	tttgggtttt	tgttattttt	1020
ttgttttagt	atttttgagt	agttgggatt	atgggtgttg	ttattgtgtt	tggttaattt	1080
tttgattttt	ttagtagaga	tggggtttta	ttgtgttagt	taggatgggt	ttgatttttt	1140
gatttttgtga	tttattttatt	ttgggttttt	aaagtgttgg	gattataggt	gtgagttatt	1200
gtgtttgggt	tgtttttgtt	tttttttttt	ttgagatagt	tttatttttg	ttgttttagt	1260
tggagtgtag	tggtgtaatt	ttgggttatg	taatttttgt	tttttgggta	taagtgattt	1320
ttttgtttta	gttttttttag	tagttgtgat	tataggtatg	tattattatg	tttgggtaat	1380
tttgattttt	tagtagagat	gggggttttt	tatgttgggt	aggttgggtt	ttaatgtttg	1440
atttttaggtg	atttgttttat	tttagttttt	taaagttttg	tgattatagg	tggtgagttat	1500
tatgtttggg	tagttttttg	tttttttaaa	aattttattat	ttagttgggt	tggtgtttta	1560
tatttgtaat	tttgggtattt	tgggaggttg	aggtgagttg	attatttgag	gttaggtgtt	1620
ttagattagt	ttgggtaata	tggtaaaatt	tggtttttat	taaaaagata	aaaattagtt	1680
ggaggtgttt	ttttgaattt	aggaggtaga	ggttgtagt	aattgagatt	atgttattgt	1740
attttaattt	aggaggagga	ggttgtattg	agttaatatt	atgttattat	atttttagttt	1800
gggtgataga	gtaagatttt	gtttttttta	aaaaaaaaat	agaaaaaaaa	aaaaagaaag	1860
aaatttgaag	taattatgat	taaatgttaa	tatttgtttt	tttatgtggt	gggtataggt	1920
gtgttttggg	agttttttatg	tttttttgtat	tttgagatta	tataaaatta	tgttttattat	1980
tttatttgtg	tatatgattt	ttgtagaaaa	gttggtaaat	tttggaaatgt	atgaagaaaa	2040
agattataat	agttagtaat	atttttgttt	agagagatag	tattttattaa	tttttttttg	2100
tttatgtttg	tttttggaaa	aatgggatat	tattgggtatt	attttataat	tttttaattt	2160
gtattaatat	tttttatatt	ttttaaattg	aaaaagtgtg	ttaagtatgg	tggttttaggt	2220
ttgtaaattt	agttaatgtg	ggaggattgt	ttgagtttgg	gagtttagga	atagtttggg	2280
tattgtgggtg	agatttatatt	gttataaaa				2308

<210> 26

<211> 2308

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 26

ttttagtagtaa	tgtgggttttg	ttatgatggt	taggttgttt	ttgaattttt	gggtttaagt	60
gatttttttta	tattgggttgg	gtttataggt	ttgagttatt	atgttttggt	agttttttta	120
atttaaaaaa	tatagaaagt	gttaataata	attaaaaagt	tataaaatag	tgtaaatagt	180
attttatttt	tttgaagata	aatataggta	gagaaaaatt	aatggatatt	gttttttttg	240
gtaggggtat	tattggttat	tataattttt	tttttttgtg	attttagaat	ttattaattt	300
ttttataaag	attatgtagt	ataataagat	aataaatata	attttgtgtg	attttaaaat	360
ataggaagta	tagaagtttt	taaaaatagt	ttgtatttat	tatgtgaaga	aatagatggt	420
aatattttgg	tataattggt	ttagattttt	tttttttttt	ttttttttgt	tttttttttg	480
gggggggatg	agtttttgtt	tggtgttttag	gttggaaatgt	agtggatatga	tattggttta	540
gtgtataattt	tttttttttg	gttggagttg	agtggtagta	tttttagttta	ttgtaatttt	600
tggttttttg	gtttaagaga	gtatttttag	ttaatttttg	tttttttagt	agagataagg	660
ttttgttatg	ttgttttagt	tggttttgaa	tatttgattt	taagtgattt	gtttattttg	720
gttttttaaa	gtgttaggat	tataggtgtg	agatattata	ttggttaaat	aataagtttt	780
taaaaaaata	ggagattggt	taggtatggt	gattttatgt	tgtaatttga	gaattttggg	840
agggttaggt	gggtagatta	tttgagattg	ggtagttggg	attagtttgg	ttaatatgga	900
gaaattttgt	ttttattaaa	aatataaaaat	tagttgggtg	tggtgggtgtg	tgtttgaat	960
tatagttatt	agggaggttg	aggtaggaga	attgtttgta	tttgggaggt	ggaggttgtg	1020
tgagttaaga	ttgtgttatt	gtatttttagt	ttgggtaatt	agaataaaaat	tgttttaaaa	1080
aaaaaaaaaa	taaaaatagg	ttgggtgtgg	tggtttatgt	ttgtaatttt	agtatttttg	1140
aaggttaagg	tgggtggatt	atgaggttag	gagattgaaa	ttatttttgt	taatatggtg	1200
aaattttgtt	tttattaaaa	attagttggg	tggtgtgggt	gtgtttgttag	gtgtttgttag	1260
tttttagttat	ttgggggtgt	tgaggttagga	gaatgggtgag	aatgtgggag	gtagagtttg	1320
tagtgagttg	agattgtgtt	attgttagttt	gggttgggtg	aaagagtgag	attttgtttt	1380
aaaaaaaaaa	aaattaaata	aataaaataaa	taaaaaaata	aaataatagg	agaagatgat	1440
gatagatttt	ataagagttt	gggttttagg	ttgaaatttt	atttagtgtt	gtggttttgt	1500
gggtgtgggt	ttatgggtgt	ttttaaaggg	tttttaggtag	atggaggttt	ttgggtgggtg	1560
tggaggaagg	aaaaggatga	aattgggtttt	tttgggttatt	tttttttagtt	tagttgtagt	1620
tgtagtttgt	gggttttagt	tgattttttg	gggtttattg	tttattttat	agatgtggag	1680
tttatagttg	ggaggtgtgg	tgggttttgg	tggtgtaggg	ttttgggggtg	gggaggttga	1740
ggtgttttgt	tttttttgtt	gggttttatt	gggttagttt	tgagtttttg	agtagggtgg	1800
attggttgtt	gtttgtttaa	agttgggggt	gttaggtagt	tgattgtttg	tggattttga	1860
gttgggtatt	gggtgtaatg	tttgttttgt	gatggttaggt	gagatgtggg	tggttttgtg	1920
ggaggggttt	ttgtttgttt	ttgttattta	tttttttttt	gtttgttgtt	gttgtagggg	1980
tgggaatggg	ggggattaag	tattttgttag	gggtgatttg	ggagattttg	gagtttagtg	2040
tttttagtgtt	gaatgagagt	tttttttttg	gtttgtgttt	gtagggattg	aagggatttt	2100

tgaatgtttg	tgggatgtgt	tgtggtggtg	tttagtatgg	aggaggtagg	tagggaggtg	2160
aggagtaggt	tagattgggt	gtagggttag	ggttttttgt	tttttaggt	tgtggttttg	2220
tggagggttt	gtagatagtt	gttagatta	tgagggttag	aggtttgttg	ttatgtgtgt	2280
tattattagt	tagttgtttt	ttgtttgg				2308

<210> 27
 <211> 2553
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 27

ttggttattt	ttgttttttt	ttagattttt	agagttttta	gtgtagttat	agaggatggt	60
gttggtttta	gttttaagaa	gatgttgttt	tttttagagg	gttaagtaag	tgggaatttt	120
tttttttatt	tgttttgggt	tttaggtagg	gtttttgggt	taaggtttgg	ggttggaagt	180
tgatttattt	aggttttaggt	tttggggtag	aattgaaatt	ttttggttat	tgttgggtgt	240
agtttgggag	taggttattg	ttaaagtgt	gggttttttt	aggatagttt	ttttatgagg	300
ttggtttttt	atttgttgtt	ttttatatatt	tggtgggttag	ggatgtggtt	ttgggtagaa	360
tgatgatttt	ttatttttgt	tattatggaa	gttatgtgtg	tttttagttt	tagtttagtta	420
tttgggttgt	agagtatttt	ttttatgttt	tttgggtgtt	tttttttttt	tttgttttag	480
tttgggtttg	ttttattttg	tttttaggga	gggttatttt	ggagtggggt	tagggtaggg	540
tttttttttg	agggagtttt	tttttgggtt	tttttagggg	agttttgtat	agtttttagta	600
tttgggtgat	tttttttgat	atttttttta	gggtagttaa	ggtattttgt	gtgggggtatt	660
taagagagtt	aggtttgtta	gttttttagtt	tttgtagtaa	tgtaggtttg	aggggtgagg	720
ggtagggtag	ggtaggggat	aggaattttg	gtgtgttttt	tatttgtaaa	ggtttattga	780
gggttttagt	tttagttatt	gagttattaa	gttagtttgg	gttaggtttg	ggtgttttgt	840
ttgtaattga	ggtagagatg	gggttttggg	gtagttttga	ggatgttggg	tgtagtagtg	900
gggttttgtt	ggtaggaatt	atttatgttt	tttttgggtt	taagttttgt	ggatgttttag	960
tttgggggtg	tggtgggttg	gtagggttagt	ggtagatatt	ggtaggttaga	tttaggtttt	1020
ggtagaatag	gtatttaagga	agtgggtgatt	ggagggaagt	taagtgtatt	taaattttttg	1080
ggtagagttat	tattgtttggg	tttttttatag	ttgttgaaag	tgagtaatag	tgatgaaggt	1140
ttgtgagttt	ttgtgtgagt	gagtgaatgg	attagtagta	gttttttaggt	tgtaggaagag	1200
tgtttttttt	ttgggatggg	gatattttgt	tatagtaatt	tttaattttt	tattttattta	1260
ttgtttattg	tagaggtagt	tggtgggtttt	gttttttgta	ggtaggagtg	aggggtattt	1320
ttgtgatgtg	gtatttttgt	gattgagggt	atgtgtgatt	gggtgaaggg	taggaagtga	1380
gttattgggt	tgtattaggt	gtgggggttt	ttgtgagggt	aggatttaaa	gttgggttgg	1440
tttttttggt	gtagtatttt	tttttttttt	gaattaggtt	agagttttgg	gatgggaggt	1500
gttttgtaga	ttattttttt	tatttaatttt	tgttttttgt	tttatttttg	tggtgatttg	1560
gtgaatttgt	ggttttttgt	tgtgtattga	gtggggtagt	gatttttgat	tgggtgtttt	1620
tggtgttttt	gttattgtta	ttatttttgg	tggttttagtt	tttgtatttt	ttatttttat	1680
ggaggaatgt	attaggtttt	tttttttggg	tgtatttttt	atttatatgt	ttttaaat	1740
tggtattttt	tgtttttttt	ttatttttat	tttttttttt	aggtttttag	ataaagggga	1800
agtggtttga	tttttttaaa	gggtagtgtt	tttattagtt	tattgttgaa	tttttttttt	1860
taatttttagt	tttttagtta	tagttaatta	gtatttagtag	atagtttatg	agtgatattt	1920
atgtaggttt	taggttgtgg	agagtttttt	gggtaggaaa	tagtttttaa	ggttttttat	1980
tttatttagg	tttttagttt	ttttatttgt	ttttttttta	gattgtggtt	ttttggagtt	2040
tggttttttt	gttttttgtt	gattgatata	tagtatttaa	atagtggtag	agtgggatgg	2100
attttttagt	ttgttttttg	tgtgggtttg	tatttttagt	tagatatgtt	tttttagatt	2160
aggatttagg	gggtatatag	tgtgtttgtg	gggtttattg	gggtatttga	tttgggttat	2220
tttatttttt	agagagaggg	ttttgttgtg	ttatttagtt	ggagtgtagt	ggtgttaatta	2280
tagtatattg	tagtttttaa	tttttgggtt	taagtgattt	ttttttttta	gttttttttag	2340
tagttgggag	tataggattt	attgtatttt	ggttaatttt	tttaataattt	tttaagagat	2400
gggggtttat	tgtgttgttt	agggttggtt	ttaaattttt	gttttaagt	atttttttat	2460
ttttgttttt	tgaagtgttg	agattatagg	tatgagttat	tatgtttatt	ttagattgat	2520
atttttatat	ttgtttattt	tggttgggta	ggg			2553

<210> 28
 <211> 2553
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 28

ttttgttttag	ttaggatgaa	taaatataga	aatgttagtt	tgggatgggt	atgggtggtt	60
atgtttgtaa	tttttagtatt	ttaggagggtg	aagggtgggag	gattatttga	ggtttagaggt	120

ttgaggtag	tttgggtaat	atagtaagat	tttatttttt	aaaaaattat	taaaaaatta	180
gttaggatat	agtgggtttt	gtattttttag	ttattaggga	ggttggggag	ggaggattgt	240
ttgagtttag	gagttgaagg	ttgtagtgtg	ttatgattat	attattgtat	tttagttggg	300
tgatatagta	agattttttt	tttaaaaaat	aaaataaatt	aaatgtgggt	gttttagtga	360
atttgttaggt	atatatgtgt	ttttttgggt	tttgttgtgg	gggggtatgt	ttgggttagg	420
gtatagattt	atatagagaa	taggttaggg	ggtttgtttt	gttttgttat	tgtttgggtg	480
ttatgtgttg	gttatatagg	gatagaagaa	ttaggtttta	aagggttata	atttaggaga	540
gaggtaggta	ggaaagattg	ggatttggat	gagatgagg	attttaaggg	ttgtttttta	600
tttaggaaat	tttttatagt	ttgggggttg	tatgggtatt	atttatgggt	tgtttgttaa	660
tgtaatttaa	ttgtaattag	ggaattgggg	ttgaggaggg	gagtttagta	gtaagtgggt	720
gggatatatt	ttttttaaga	ggatttaggt	attttttttt	tgtttgggag	tttaggggaa	780
gggggtgggag	taaaggggga	gtagaaaatg	ttagggtttg	gaagtatgtg	ggtgaggggt	840
gtatttagga	agggaggttt	ggtgtatttt	tttatggggg	tggggaatgt	ggaggttggg	900
ttattggagg	tggtgggtgg	ggtagggggt	gtaggagatg	ttatgttagg	gttattgttt	960
tatttgggtg	atagtgggg	gttggtagtt	tattggatta	ttgtgggggt	gggggtggagg	1020
atggaagtgt	gtgagggggg	tggtttatag	ggtatttttt	gttttagggg	tttaatttaa	1080
tttgaaaggg	aaaggagtgt	tgtagttggg	aggttagggt	gattttgggt	tttgtttttg	1140
tagaagtttt	tatgtttggt	gtagattaat	gatttgtttt	ttgtttttat	attgattata	1200
tatgattttg	gttatagggg	tgttatatta	taggagtgtt	ttttattttt	gtttgttaga	1260
agtagggttt	ttgtatattt	ttgtagtggg	tggtgggtgg	gtgggggatt	aggaattggt	1320
gtaattaaagt	gtttttattt	tggggagggg	atgttttttt	ataatttggg	aattgttatt	1380
ggttttattt	tttgtttatg	tagaaattta	taaattttta	ttattgttgt	ttatttttag	1440
tagttgtgaa	agatttgggt	gtgatgattt	atttgagggg	ttgagtgtat	ttggtttttt	1500
tttgggttatt	atttttttga	tgtttgtttt	atttagattt	aggtttgttt	attagtgttt	1560
gttattgatt	tgttagtttt	ttgtgggttt	aggttgggta	tttataaagt	ttggtttagg	1620
agagagtata	agtgattttt	gttgggtagg	tttttgttgt	gtatttagta	tttttagaat	1680
tgttttgaga	ttttgttttt	gtttttattg	tagatagggg	atttaggttt	ggtttaggtt	1740
gatttgggtg	tttagtggtt	ggggtttggg	gttttagtga	atttttgtgg	atggagagta	1800
tgttggaggt	tttgtttttg	tttttgtttt	gttttttatt	tttttaggtt	gtattttggg	1860
aggagttaga	ggttgatagg	tttgggtttt	ttgagtgttt	tatatagagt	gtttgattgt	1920
ttttaagtaag	gatgttaagg	gaggtgtgtt	aggatttgag	gttgtgtaga	gttgttttgg	1980
ggatagttag	agaggaattt	ttttggggga	gagttatgtt	ttggttttat	tttaggggat	2040
ttttttttga	gagtagggta	ggataaagtt	aggttggggg	aggagaaggg	ggaggatttt	2100
ggaggggtatg	aaaggggtgt	tttgtagttt	agggtggttg	ttgggttggg	agatagtggg	2160
ggtttttata	atgataggag	tggagaatta	ttgttttatt	taggggtata	tttttggtta	2220
ttagggtgtga	agaaatagta	ggtggaggat	tggtttttatg	gggagattgt	tttggaagga	2280
tttatagttt	tggtagtggg	ttgttttttag	gttgtagtgt	atagtaatta	aggagtttta	2340
gttttgtttt	agagtttgga	tttaggtggg	ttggtttttta	gttttaggtt	ttatattagg	2400
ggttttgttt	ggagttttag	ataagtaggg	agggggattt	ttattttatt	agttttttgg	2460
aggaagtggg	gttttttttg	ggttgaagtt	aatagtattt	tttgtgggta	tattgggggt	2520
tttgagggtt	taagggaggg	taggggtggg	tag			2553

<210> 29
 <211> 2308
 <212> DNA
 <213> Homo Sapiens

<400> 29

ccaggcagga	ggcagctagc	tggtgatagc	gcgcgtgacc	acagacctct	gacctcata	60
gtctcagcag	ctgtctgcag	gccctccacg	ggaccgcagc	ctgggagggc	aggaggccct	120
gcacctgcgg	cgggtctgac	ctgctcctca	cctccctgcc	tgccctctcc	gtgctgggcg	180
ccgccgcgac	acgtcccgcg	gacattcagg	agtccttcca	gtccctgcga	gcgcggaccc	240
gggagagagc	tctcattcaa	cgtgaggcc	actggctcac	agatctcccc	gggtcaccct	300
gcgaagtgct	tggtcccccc	cattcccagc	cctgcagcag	caacaaacgg	gaaggaagtg	360
ggtgacgagg	acaggcagag	acccctccca	cggggccgcc	cgcactctac	ctgccgtcgc	420
ggggcgggcg	ttgcggcgcg	tgcccagctg	cagatccgca	ggcgatcgca	ctcctggcgg	480
ccccagcttt	aagcaggcag	cggccgatcc	gccctgctcg	agggtctcagg	gctgaccag	540
gtgagccccg	gcggggaggg	cgggcgcctc	ggcctccccg	ccccggaaac	ctgcaccgcc	600
cagagccgcc	acgcctcccc	gctgtaggct	ccgcactctg	aaaatggggc	gatgaacccc	660
aagggtcggc	tcagacccac	ggactgcagc	tgcagctggg	ctagaagggg	tggccagggg	720
agccgatttc	atcccttttc	tccctccacg	ccaaccagga	gcctccatct	gcctgggacc	780
ctttgggggc	acccatgggc	ccacacgcg	agaaccacag	cactgggtgg	ggtttcagcc	840
ctggggccaa	gctcttgtga	ggtctgtcat	catcttctcc	tgttgttttg	tttttttgtt	900
tgtttgtttg	tttggttttt	ttttttttga	gacagagtct	cgctctttca	cccaggccgg	960
actgcagtgg	cgcgatctcg	gctcactgca	agctctgcct	cccgggttct	cgccattctc	1020
ctgcctcagc	acccccagag	agctgggact	acgggcgcgc	ccaccgcgcc	cggctaattt	1080
tttgtatttt	ttagtagaga	cggggtttca	ccgtgttagc	caggatgggt	tcgatctcct	1140
gacctcgtga	tcacccacc	ttggccttcc	aaagtgctgg	gattacaggc	gtgagccacc	1200
gcgcccggcc	tgtttttgtt	tttttttttt	ttgagacagt	tttattctgg	tgcccaggc	1260
tggagtgcag	tggcgcaatc	ttggctcacg	caacctccgc	ctcccgggta	caagcgattc	1320

tcctgcctca	gcctccctag	tagctgtgat	tacaggcacg	caccaccacg	cccggcta	1380
tttgtatttt	tagtagagac	ggggtttctc	catgttgcc	aggctggtct	ccaatgccc	1440
atctcaggtg	atctgcccac	ctcagcctcc	caaagtctcg	cgattacagg	cgtaggtcac	1500
catgcctggc	cagtctcctg	tttttttaaa	aacttattat	ttagccggtg	tggtgtctca	1560
cacctgta	cctggcactt	tgggaggccg	aggtagcg	atcacttgag	gtcagggtgt	1620
ccagaccagc	ctgggcaaca	tggcaaaacc	ttgtctctac	taaaaagaca	aaaattagct	1680
ggaggtgctc	tcttgaaccc	aggaggcaga	ggttgcagtg	aactgagatc	atgccactgc	1740
actccaaccc	aggaggagga	ggttgcactg	agccaatata	atgccactac	attccagcct	1800
gggcgacaga	gcaagactct	gtccccccca	aaaaaaaaaac	agaaaaaaaaa	aaaaagaaag	1860
aaatctgaag	caattatgac	caaagtgtta	catctgtttc	ttcacgtggt	gggtacaggc	1920
gtgttttggg	agcttctatg	cttctctgat	tttgagatca	cacaaaatta	tgttttattat	1980
cttattgtgc	tacatgatct	ttgtagaaaa	gttggtaaat	tctggaatgc	acgaagaaaa	2040
agattataat	agccagta	acccctgccc	agagagacag	tatccattaa	tttttctctg	2100
cctatgtttg	tcttcggaaa	aatgggatac	tattggcact	attttataac	tttttaattt	2160
gtattaacac	tttctatatt	ttttaaattg	aaaaagctgg	ccaagcatgg	tggctcaggt	2220
ctgtaaacc	agccaatgtg	ggaggatcgc	ttgagcccg	gagttcagga	acagcctggg	2280
catcgtggcg	agaccacatt	gctacaaa				2308

<210> 30
 <211> 2553
 <212> DNA
 <213> Homo Sapiens

<400> 30

ctggccaccc	ctgccctccc	ttagacctcc	agagccccc	gtgtagccac	agaggatgct	60
gttggttcca	gccccaaaga	gacgccgctt	cctccagagg	gctaagtaag	tggaatccc	120
cctccctact	tgctctgggc	tccaggcagg	gccccgtgtg	taaggcctgg	ggctggaagc	180
cgaccacact	aggtccaggc	tctggggcag	aactgaaact	ccttggttac	tgctggctgc	240
agcctgggag	caggccactg	ccaaagctgt	gggtccctcc	aggacagtct	ccccatgagg	300
ccgggtctct	acctgctgtt	tcttcacacc	tggtggccag	ggatgtggcc	ctgggtagaa	360
cgatgattct	ccactcctgt	cattatggaa	gccaccgtcg	tctccagacc	cagccagcca	420
cctgggctgc	agagcacc	tttcatgccc	tccgggtgcc	tcccccttct	cctgccccag	480
cctggctttg	tcctaccctg	ctctcaggga	gggtgacct	ggagtggggc	cagggcatgg	540
ctctcccccg	agggagtccc	tctctggctg	tccccagggc	agctctgcac	agcctcagta	600
cctggcgcac	ctcccttgac	atccttctta	gggacagtca	ggcactctgt	gtggggcact	660
caagagagcc	agggccgtca	gcctctagct	cctgccagaa	tgcaggcctg	aggggtgagg	720
ggcggggcag	gggcaggggc	aggaactccg	gcgtgctctc	catccgcaaa	ggttcactga	780
ggccccgagc	cccagccact	gagccaccaa	gtcagcctgg	gccaggcctg	gggtgccctgt	840
ctgcaatgga	ggcagagacg	gggtctcggg	gcagtctctga	ggatgctggg	tgcacagcgg	900
gggcctcgcc	ggcaggaatc	acttatgctc	tctcctgggc	caagctttgt	ggatgcccag	960
cctggggccg	cggggagctg	gcaggctcagt	ggcagacact	gggtgggcaga	cctagtgtct	1020
ggtagaacag	gcataagga	agtgggtgacc	ggagggaagc	caagtgcact	caaaccctcg	1080
ggtagtcat	caccgcccgg	tctttcacag	ctgctgaaag	tgagcaacag	tgatgaaggt	1140
ttgtgagttt	ctgcgtgagc	gagtgaatgg	accagtagca	gtttccagggt	tggtgaagag	1200
cgttccctcc	ccgggatggg	gacacttggt	tacagcaatt	cctaattccc	caccaccca	1260
ccgcccactg	cagaggatag	cgggggccct	gcttctgca	ggcaggagt	aggggcactc	1320
ctgtgatgtg	gcaccctgt	gaccgaggtc	atgtgtgac	gggtgaagg	caggaagcga	1380
gtcattggtc	tgcaccaggc	gtgggggctt	ctgcgagggc	aggacccaaa	gtcggcctgg	1440
cctcccggct	gcagcactcc	tttccctttc	gaattagggt	agagccctgg	gacgggaggt	1500
gccctgtaga	ccacccccct	caccaacttc	cgctctccgc	cccacccccg	cgggtgatccg	1560
gtgaactgcc	ggccccctgc	tgtgcaccga	gtggggcagt	gacctgaag	tggcgctctc	1620
tgccgcccc	gccaccgcca	ccacctccgg	tggcccagcc	tccgcattcc	ccacccccat	1680
ggaggaatgc	accaggcctc	ccttctctgga	tgcacccctc	acccacatgc	ttccaaaccc	1740
tggcattttc	tgctccccct	ttactcccac	cccttcccct	aggctcccag	acaaagggga	1800
agtggctgga	tcctcttaaa	gggacagtgt	cccaccagct	tactgctgaa	ctccccctct	1860
caacccagct	tccttagtta	cagtttaatta	gcattagcag	acagcccattg	agtgaatacc	1920
atgcaggccc	caggctgtgg	agagtttctt	gggtaggaaa	cagcccttaa	gggtccctcat	1980
ctcatccagg	tcctagttct	tcctacctgc	ctctctctta	gattgtggcc	ctttggagcc	2040
tggttcttct	gtccctgtgt	gaccgacaca	tagcacccaa	acagtggcag	agcgggacgg	2100
accccttagc	ctgtttctctg	tgtgggtctg	taccctgacc	cagacatgcc	ccccacagc	2160
aggacccagg	ggggcacatg	tgtgcctg	ggttcactgg	ggcaccgcga	tttggtttat	2220
tttatttttt	agagagaggg	tcttgtctg	tcaccagct	ggagtgcagt	gggtgaatca	2280
tagcacactg	cagccttcaa	ctcctgggct	caagcgatcc	tcctcccca	gcctccctag	2340
tagctgggag	tacaggaccc	actgtatcct	ggctaatttt	ttaataattt	tttaagagat	2400
ggggtcttac	tgtgttgccc	aggctggcct	caaacctctg	gcctcaagt	atcctcccac	2460
cttcgcctcc	tgaagtgtctg	agattacagg	catgagccac	catgcccatc	ccagactgac	2520
atttctatat	ttgttcatcc	tggctgggca	ggg			2553

<210> 31

<211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)

 <400> 31
 gggattattt ttataagggt 20

 <210> 32
 <211> 17
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)

 <400> 32
 ctctaaaccc catcccc 17

 <210> 33
 <211> 24
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)

 <400> 33
 cccatcccca aaaacacaaa ccac 24

 <210> 34
 <211> 19
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)

 <400> 34
 cgtcgctgta gttttcggt 19

 <210> 35
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)

 <400> 35
 tagtgagtac gcgcggtt 18

 <210> 36
 <211> 19
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)

 <400> 36
 accgaaaata cgcttcacg 19

<210> 37
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)
 <400> 37
 gcgttatcgt aaagtattgc gc 22

 <210> 38
 <211> 19
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)
 <400> 38
 cgcgacgaac aaaacgccg 19

 <210> 39
 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)
 <400> 39
 cgcgctactc cgcataca 18

 <210> 40
 <211> 19
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)
 <400> 40
 gaggtaatcg aggcggtcg 19

 <210> 41
 <211> 21
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)
 <400> 41
 cgccaattca tacgccgcac c 21

 <210> 42
 <211> 19
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> chemically treated genomic DNA (Homo sapiens)
 <400> 42
 ttgtggttcg ggaagagac 19

<210> 43
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)
<400> 43

tcccgaactc ttcgatcg 18

<210> 44
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)
<400> 44

aactacgcg aaacccgcga 20

<210> 45
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)
<400> 45

cgcgctactc cgcataca 18

<210> 46
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)
<400> 46

gaggtaatcg aggcggtcg 19

<210> 47
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)
<400> 47

cgccaattca tacgccgcac c 21

<210> 48
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)
<400> 48

accgaaaata cgcttcacg	19
<210> 49	
<211> 22	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> chemically treated genomic DNA (Homo sapiens)	
<400> 49	
gcggttatcgt aaagtattgc gc	22
<210> 50	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> chemically treated genomic DNA (Homo sapiens)	
<400> 50	
cgcgacgaac aaaacgccg	19
<210> 51	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> chemically treated genomic DNA (Homo sapiens)	
<400> 51	
gcgtttttacg tcgtcgcg	18
<210> 52	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> chemically treated genomic DNA (Homo sapiens)	
<400> 52	
gacgctaaac gccaccgt	18
<210> 53	
<211> 23	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> chemically treated genomic DNA (Homo sapiens)	
<400> 53	
ccgaccatcc gacgccttac tcg	23
<210> 54	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> chemically treated genomic DNA (Homo sapiens)	
<400> 54	

cgtttttcgt tttatttcg c 21

<210> 55
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 55

gacaaaaaac gccacgtc 18

<210> 56
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 56

ccgacaattc accgaatcac cg 22

<210> 57
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 57

atctcaccta ccgtcgcg 18

<210> 58
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 58

taggagtgcg atcgtttgc 19

<210> 59
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> chemically treated genomic DNA (Homo sapiens)

<400> 59

acgaacgtta cgaccgatac ccaacta 27